

Methods of Optimizing Substrate Pools and Biosynthesis of
Poly- β -Hydroxybutyrate-co-Poly- β -Hydroxyvalerate
in Bacteria and Plants

Background of the Invention

Field of the Invention

The present invention relates to genetically engineered plants and
10 bacteria. In particular, it relates to methods for optimizing substrate pools
to facilitate the biosynthetic production of commercially useful levels of
polyhydroxyalkanoates (PHAs) in bacteria and plants.

PHAs are bacterial polyesters that accumulate in a wide variety of
15 bacteria. These polymers have properties ranging from stiff and brittle
plastics to rubber-like materials, and are biodegradable. Because of these
properties, PHAs are an attractive source of nonpolluting plastics and
elastomers.

20 The present invention especially relates to the production of
copolyesters of β -hydroxybutyrate (3HB) and β -hydroxyvalerate (3HV),
designated P(3HB-co-3HV) copolymer, and derivatives thereof.

Description of Related Art

25

Currently, there are approximately a dozen biodegradable plastics
in commercial use that possess properties suitable for producing a number
of specialty and commodity products (Lindsay, 1992). One such
biodegradable plastic in the polyhydroxyalkanoate (PHA) family that is
30 commercially important is Biopol™, a random copolymer of 3-
hydroxybutyrate (3HB) and 3-hydroxyvalerate (3HV). This bioplastic is

used to produce biodegradable molded material (e.g., bottles), films, coatings, and in drug release applications. Biopol™ is produced via a fermentation process employing the bacterium *Alcaligenes eutrophus* (Byrom, 1987). The current market price is \$6-7/lb, and the annual
5 production is 1,000 tons. By best estimates, this price can be reduced only about 2-fold via fermentation (Poirier et al., 1995). Competitive synthetic plastics such as polypropylene and polyethylene cost about 35-45¢/lb (Layman, 1994). The annual global demand for polyethylene alone is about 37 million metric tons (Poirier et al., 1995). It is therefore likely that the
10 cost of producing P(3HB-co-3HV) by microbial fermentation will restrict its use to low-volume specialty applications.

Nakamura et al. (1992) reported using threonine (20g/L) as the sole carbon source for the production of P(3HB-co-3HV) copolymer in *A.*
15 *eutrophus*. These workers initially suggested that the copolymer might form via the degradation of threonine by threonine deaminase, with conversion of the resultant α -ketobutyrate to propionyl-CoA. However, they ultimately concluded that threonine was utilized directly, without breaking carbon-carbon bonds, to form valeryl-CoA as the 3HV precursor.
20 The nature of this chemical conversion was not described, but since the breaking of carbon-carbon bonds was not postulated to occur, the pathway could not involve threonine deaminase in conjunction with an α -ketoacid decarboxylating step to form propionate or propionyl-CoA. In the experiments of Nakamura et al., the PHA polymer content was very low (<
25 6% of dry cell weight). This result, in conjunction with the expense of feeding bacteria threonine, makes their approach impractical for the commercial production of P(3HB-co-3HV) copolymer.

Yoon et al. (1995) have shown that growth of *Alcaligenes* sp. SH-69
30 on a medium supplemented with threonine, isoleucine, or valine resulted in significant increases in the 3HV fraction of the P(3HB-co-3HV) copolymer.

In addition to these amino acids, glucose (3% wt/vol) was also added to the growth media. In contrast to the results obtained by Nakamura et al. (1992), growth of *A. eutrophus* under the conditions described by Yoon et al. (1995) did not result in the production of P(3HB-co-3HV) copolymer when
5 the medium was supplemented with threonine as the sole carbon source. From their results, Yoon et al. (1995) implied that the synthetic pathway for the 3HV component in P(3HB-co-3HV) copolymer is likely the same as that described in WO 91/18995 and Steinbüchel and Pieper (1992). This postulated synthetic pathway involves the degradation of isoleucine to
10 propionyl-CoA (Figure 3).

The PHB Biosynthetic Pathway

Polyhydroxybutyrate (PHB) was first discovered in 1926 as a
15 constituent of the bacterium *Bacillus megaterium* (Lemoigne, 1926). Since then, PHAs such as PHB have been found in more than 90 different genera of gram-negative and gram-positive bacteria (Steinbüchel, 1991). These microorganisms produce PHAs using R- β -hydroxyacyl-CoAs as the direct metabolic substrate for a PHA synthase, and produce polymers of R-(3)-
20 hydroxyalkanoates having chain lengths ranging from C3-C14 (Steinbüchel and Valentin, 1995).

To date, the best understood biochemical pathway for PHB production is that found in the bacterium *Alcaligenes eutrophus* (Dawes and
25 Senior, 1973; Slater et al., 1988; Schubert et al., 1988; Peoples and Sinskey, 1989a and 1989b). This pathway, which is also utilized by other microorganisms, is summarized in Figure 1. In this organism, an operon encoding three gene products, i.e., PHB synthase, β -ketothiolase, and acetoacetyl-CoA reductase, encoded by the *phbC*, *phbA*, and *phbB* genes,
30 respectively, are required to produce the PHA homopolymer R-polyhydroxybutyrate (PHB).

As further shown in Figure 1, acetyl-CoA is the starting substrate employed in the biosynthetic pathway. This metabolite is naturally available for PHB production in plants and bacteria when these organisms are genetically manipulated to produce the PHB polyester.

5

Recently, a multi-enzyme pathway was successfully introduced into plants for the generation of polyhydroxybutyrate (PHB) (Poirier et al., 1992).

10 The P(3HB-co-3HV) Copolymer Biosynthetic Pathway

As noted above, P(3HB-co-3HV) random copolymer, commercially known as Biopol™, is produced by fermentation employing *A. eutrophus*. A proposed biosynthetic pathway for P(3HB-co-3HV) copolymer production is shown in Figure 2. Production of this polymer in plants has not yet been demonstrated.

The successful production of P(3HB-co-3HV) copolymer in plants or bacteria requires the generation of substrates that can be utilized by the PHA biosynthetic enzymes. For the 3HB component of the polymer, the substrate naturally exists in plants in sufficient concentration in the form of acetyl-CoA (Nawrath et al., 1994). This is not true for the 3HV component of the copolymer, however. In this case, the starting substrate is propionyl-CoA. The presence of sufficient pools of acetyl-CoA and propionyl-CoA in plants and microorganisms, along with the proper PHA biosynthetic enzymes (i.e., a β -ketothiolase, a β -ketoacyl-CoA reductase, and a PHA synthase), would make it possible to produce copolyesters of P(3HB-co-3HV) in these organisms.

Summary of the Invention

The present invention provides a variety of different methods for optimizing levels of substrates employed in the biosynthesis of copolymers of 3-hydroxybutyrate (3HB) and 3-hydroxyvalerate (3HV) in plants and bacteria via manipulation of normal metabolic pathways using recombinant DNA techniques.

In one aspect, the present invention provides methods for the production of enhanced levels of threonine, α -ketobutyrate, propionyl-CoA, β -ketovaleryl-CoA, and β -hydroxyvaleryl-CoA, all of which are metabolites in the biosynthetic pathway of 3HV in the P(3HB-co-3HV) copolymer, from various carbon sources in plants or bacteria. This is achieved by providing a variety of wild-type and/or deregulated enzymes involved in the biosynthesis of the aspartate family of amino acids (i.e., aspartate, threonine, lysine, and methionine), and wild-type or deregulated forms of enzymes involved in the conversion of threonine to P(3HB-co-3HV) copolymer endproduct. Using these enzymes, the levels of the above-noted metabolites can be increased in plants and bacteria in the range of from about 1-10 fold, 1-100 fold, or 1-1000 fold.

In another aspect, the present invention provides methods for the biological production of P(3HB-co-3HV) copolymers in plants and bacteria utilizing propionyl-CoA as a substrate. As shown in Figure 3, propionyl-CoA can be produced through a variety of engineered metabolic pathways. Introduction into plants and bacteria of appropriate β -ketothiolases capable of condensing acetyl-CoA with itself and/or with propionyl-CoA, along with appropriate β -ketoacyl-CoA reductases and PHA synthases, in combination with various enzymes involved in aspartate family amino acid biosynthesis and the conversion of threonine to PHA copolymer precursors,

will permit these organisms to produce P(3HB-co-3HV) copolymers. The PHA biosynthetic enzymes can be those of *A. eutrophus* or other organisms, or enzymes that catalyze reactions involved in fatty acid biosynthesis or degradation, ultimately resulting in the conversion of acetyl-CoA and propionyl-CoA to P(3HB-co-3HV). In plants, these enzymes can be expressed in the cytoplasm or targeted to organelles such as plastids (e.g., those of leaves or seeds) or mitochondria via the use of transit peptides for enhanced production of polyhydroxyalkanoates. Alternatively, plastids can be transformed with recombinant constructs that facilitate expression of these enzymes directly within the plastids themselves.

The enzymes discussed herein can be employed alone or in various combinations in order to enhance the levels of threonine, α -ketobutyrate, propionate, propionyl-CoA, β -ketovaleryl-CoA, and β -hydroxyvaleryl-CoA, and for the production of P(3HB-co-3HV) copolymer.

More specifically, the present invention encompasses the following aspects:

An isoleucine-deregulated mutein of *E. coli* threonine deaminase, wherein leucine at amino acid position 447 is replaced with an amino acid selected from the group consisting of alanine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine. In one aspect, leucine at amino acid position 447 can be replaced with phenylalanine.

An isoleucine-deregulated mutein of *E. coli* threonine deaminase, wherein leucine at amino acid position 447 is replaced with an amino acid selected from the group consisting of alanine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine, and leucine at amino acid position 481 is replaced with an amino acid selected from the group

consisting of alanine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine. In one aspect, leucine at amino acid position 447 can be replaced with phenylalanine, and leucine at amino acid position 481 can be replaced with phenylalanine.

5

An α -ketoacid dehydrogenase complex, comprising an α -ketoacid decarboxylase E1 subunit, a dihydrolipoyl transacylase E2 subunit, and a dihydrolipoyl dehydrogenase E3 subunit, wherein said α -ketoacid decarboxylase E1 subunit exhibits improved binding and decarboxylating properties with α -ketobutyrate compared to pyruvate decarboxylase E1 subunit naturally present in a host cell pyruvate dehydrogenase complex. The α -ketoacid decarboxylase E1 subunit can be a branched-chain α -ketoacid decarboxylase E1 subunit such as bovine kidney, *Pseudomonas putida*, or *Bacillus subtilis* branched-chain α -ketoacid dehydrogenase E1 subunit.

An α -ketoacid dehydrogenase complex, comprising an α -ketoacid decarboxylase E1 subunit, a dihydrolipoyl transacylase E2 subunit, and a dihydrolipoyl dehydrogenase E3 subunit, wherein said E1 and E2 subunits exhibit improved binding/decarboxylating and transacylase properties, respectively, with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit and dihydrolipoyl transacetylase E2 subunit, respectively, naturally present in a host cell pyruvate dehydrogenase complex. The α -ketoacid decarboxylase E1 and dihydrolipoyl transacylase E2 subunits can be branched-chain α -ketoacid dehydrogenase E1 and E2 subunits such as those from bovine kidney, *Pseudomonas putida*, or *Bacillus subtilis*.

An isolated β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA.

An isolated β -ketothiolase capable of condensing acetyl-CoA and
5 butyryl-CoA to produce β -ketocaproyl-CoA.

An isolated β -ketothiolase capable of:

condensing two molecules of acetyl-CoA to produce acetoacetyl-
10 CoA;

condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-
CoA; and

15 condensing acetyl-CoA and butyryl-CoA to produce β -ketocaproyl-
CoA.

The foregoing isolated β -ketothiolase is exemplified by BktB β -keto-
thiolase having the amino acid sequence shown in SEQ ID NO:11.
20

Isolated DNA molecules comprising a nucleotide sequence encoding
the isoleucine-deregulated muteins of *E. coli* threonine deaminase
described herein.

25 An isolated DNA molecule comprising the nucleotide sequence
shown in SEQ ID NO:5.

An isolated DNA molecule comprising the nucleotide sequence
shown in SEQ ID NO:8.

An isolated DNA molecule comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence shown in SEQ ID NO:9 or the
5 complement thereof;

(b) a nucleotide sequence that hybridizes to said nucleotide sequence of (a) under a wash stringency equivalent to 0.5X SSC to 2X SSC, 0.1% SDS, at 55-65°C, and which encodes an enzyme having enzymatic
10 activity similar to that of *A. eutrophus* BktB β -ketothiolase;

(c) a nucleotide sequence encoding the same genetic information as said nucleotide sequence of (a), but which is degenerate in accordance with the degeneracy of the genetic code; and
15

(d) a nucleotide sequence encoding the same genetic information as said nucleotide sequence of (b), but which is degenerate in accordance with the degeneracy of the genetic code.

20 The foregoing isolated DNA molecule, wherein said wash stringency is equivalent to 2X SSC, 0.1% SDS, at 55°C.

The foregoing isolated DNA molecule, wherein said wash stringency is equivalent to 1X SSC, 0.1% SDS, at 55°C.

25

The foregoing isolated DNA molecule, wherein said wash stringency is equivalent to 0.5X SSC, 0.1% SDS, at 55°C.

An isolated DNA molecule, comprising the nucleotide sequence
30 shown in SEQ ID NO:9 or the complement thereof.

A recombinant vector, comprising any of the foregoing nucleotide sequences operatively linked to a promoter and 5' and 3' regulatory sequences sufficient to drive expression of said nucleotide sequences in a host cell.

5

A recombinant vector, comprising a nucleotide sequence encoding *E. coli* threonine deaminase wherein leucine at position 481 is replaced with phenylalanine, operatively linked to a promoter and 5' and 3' regulatory sequences sufficient to drive expression of said nucleotide sequence in a

10 host cell.

The foregoing recombinant vector, wherein said nucleotide sequence comprises the sequence shown in SEQ ID NO:7.

15 A host cell, comprising any of the foregoing recombinant vectors.

A host cell, comprising an α -ketoacid dehydrogenase complex comprising an α -ketoacid decarboxylase E1 subunit, a dihydrolipoyl transacylase E2 subunit, and a dihydrolipoyl dehydrogenase E3 subunit, wherein said α -ketoacid decarboxylase E1 subunit exhibits improved binding and decarboxylating properties with α -ketobutyrate compared to pyruvate decarboxylase E1 subunit naturally present in a host cell pyruvate dehydrogenase complex, wherein said host cell produces an increased amount of propionyl-CoA compared to a corresponding host cell comprising wild-type pyruvate dehydrogenase complex. The α -ketoacid decarboxylase E1 subunit which exhibits improved binding and decarboxylating properties with α -ketobutyrate can be a branched-chain α -ketoacid decarboxylase E1 subunit.

20
25

A host cell, comprising an α -ketoacid dehydrogenase complex,
comprising an α -ketoacid decarboxylase E1 subunit, a dihydrolipoyl
transacylase E2 subunit, and a dihydrolipoyl dehydrogenase E3 subunit,
wherein said E1 and E2 subunits exhibit improved binding/decarboxylating
5 and transacylase properties, respectively, with α -ketobutyrate compared
with pyruvate decarboxylase E1 subunit and dihydrolipoyl transacetylase
E2 subunit, respectively, naturally present in a host cell pyruvate
dehydrogenase complex, wherein said host cell produces an increased
amount of propionyl-CoA compared to a corresponding host cell comprising
10 wild-type pyruvate dehydrogenase complex.

A plant, the genome of which comprises introduced DNA encoding a
wild-type or deregulated aspartate kinase enzyme;

15 wherein said introduced DNA is operatively linked to regulatory
signals that cause expression of said introduced DNA; and

wherein cells of said plant produce an elevated amount of threonine
compared to that in cells of a corresponding, wild-type plant not comprising
20 said introduced DNA.

A plant, the genome of which comprises introduced DNAs encoding
the following enzymes:

25 a wild-type or deregulated aspartate kinase enzyme; and

threonine synthase;

wherein said introduced DNAs are operatively linked to regulatory
30 signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an elevated amount of threonine compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

5 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase; and

10 homoserine dehydrogenase;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

15 wherein cells of said plant produce an elevated amount of threonine compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

20 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase; and

25

threonine synthase;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

30

wherein cells of said plant produce an elevated amount of threonine compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

5 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase; and

10 a wild-type or deregulated threonine deaminase;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

15 wherein cells of said plant produce an elevated amount of α -ketobutyrate compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

20 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

threonine synthase; and

25

a wild-type or deregulated threonine deaminase;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

30

wherein cells of said plant produce an elevated amount of α -ketobutyrate compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

5 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

10 homoserine dehydrogenase; and

a wild-type or deregulated threonine deaminase;

15 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of α -ketobutyrate compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

20

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

25

homoserine dehydrogenase;

threonine synthase; and

30

a wild-type or deregulated threonine deaminase;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of α -keto-
5 butyrate compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

10

a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit enzyme that exhibits improved binding and decarboxylating activities with α -ketobutyrate
15 compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex of said plant;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

20

wherein cells of said plant produce an elevated amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

25 The foregoing plant, wherein said α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate is a branched chain α -ketoacid decarboxylase E1 subunit such as that from bovine kidney, *Pseudomonas putida*, or *Bacillus subtilis*.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated threonine deaminase;

5

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant; and

10

a dihydrolipoyl transacylase E2 subunit exhibiting improved transacylase activity with α -ketobutyrate compared with dihydrolipoyl transacetylase E2 subunit naturally present in pyruvate dehydrogenase complex in said plant;

15

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an elevated amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

The foregoing plant, wherein said α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate is a branched chain α -ketoacid decarboxylase E1 subunit such as that from bovine kidney, *Pseudomonas putida*, or *Bacillus subtilis*, and said dihydrolipoyl transacylase E2 subunit exhibiting improved transacylase activity with α -ketobutyrate is a branched chain

dihydrolipoyl transacylase E2 subunit such as that from bovine kidney, *Pseudomonas putida*, or *Bacillus subtilis*.

A plant, the genome of which comprises introduced DNAs encoding
5 the following enzymes:

a wild-type or deregulated threonine deaminase; and

pyruvate oxidase;

10

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an elevated amount of propionyl-
15 CoA compared that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

20

threonine synthase;

a wild-type or deregulated threonine deaminase; and

25

pyruvate oxidase;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an elevated amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

5 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated threonine deaminase;

10 pyruvate oxidase; and

an acyl-CoA synthetase;

wherein each of said introduced DNAs is operatively linked to
15 regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an elevated amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

20

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

threonine synthase;

25

a wild-type or deregulated threonine deaminase;

pyruvate oxidase; and

30

an acyl-CoA synthetase;

Approved for Release

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an elevated amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

10

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

20

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

25

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said plant produces P(3HB-co-3HV) copolymer.

5

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

threonine synthase;

10

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

25

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said plant produces P(3HB-co-3HV) copolymer.

5

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
10 to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and
15 propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

20

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

25 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said plant produces P(3HB-co-3HV) copolymer.

The foregoing plant, the genome of which further comprises introduced DNA encoding a wild-type or deregulated threonine deaminase;

wherein said introduced DNA is operatively linked to regulatory
5 signals that cause expression of said introduced DNA; and

wherein said plant produces P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding
10 the following enzymes:

threonine synthase;

a wild-type or deregulated threonine deaminase;

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

20 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA
25 and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating

β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to
5 regulatory signals that cause expression of said introduced DNAs; and

wherein said plant produces P(3HB-co-3HV) copolymer.

Any of the foregoing plants, wherein the β -ketothiolase capable of
10 condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA is BktB.

Any of the foregoing plants, wherein said β -ketoacyl-CoA reductase is obtainable from a microorganism selected from the group consisting of
15 *Alcaligenes eutrophus*, *Alcaligenes faecalis*, *Aphanothece* sp., *Azotobacter vinelandii*, *Bacillus cereus*, *Bacillus megaterium*, *Beijerinckia indica*, *Derxia gummosa*, *Methylobacterium* sp., *Microcoleus* sp., *Nocardia corallina*, *Pseudomonas cepacia*, *Pseudomonas extorquens*, *Pseudomonas oleovorans*, *Rhodobacter sphaeroides*, *Rhodobacter capsulatus*, *Rhodospirillum rubrum*,
20 and *Thiocapsa pfennigii*.

Any of the foregoing plants, wherein said polyhydroxyalkanoate synthase is obtainable from a microorganism selected from the group consisting of *Alcaligenes eutrophus*, *Alcaligenes faecalis*, *Aphanothece* sp.,
25 *Azotobacter vinelandii*, *Bacillus cereus*, *Bacillus megaterium*, *Beijerinckia indica*, *Derxia gummosa*, *Methylobacterium* sp., *Microcoleus* sp., *Nocardia corallina*, *Pseudomonas cepacia*, *Pseudomonas extorquens*, *Pseudomonas oleovorans*, *Rhodobacter sphaeroides*, *Rhodobacter capsulatus*, *Rhodospirillum rubrum*, and *Thiocapsa pfennigii*.

A plant, the genome of which comprises introduced DNA encoding a wild-type or deregulated threonine deaminase enzyme;

wherein said introduced DNA is operatively linked to regulatory
5 signals that cause expression of said introduced DNA; and

wherein cells of said plant produce an increased amount of α -keto-butyrates compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNA.

10

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

threonine synthase; and

15

a wild-type or deregulated threonine deaminase;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

20

wherein cells of said plant produce an increased amount of α -keto-butyrates compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

25 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

30

homoserine dehydrogenase;

pyruvate oxidase;

wherein cells of said plant produce an increased amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

homoserine dehydrogenase;

threonine synthase:

20 a wild-type or deregulated threonine deaminase; and

pyruvate oxidase;

25 wherein each of said introduced DNAs is operatively linked to
regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

5

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

10

pyruvate oxidase; and

an acyl-CoA synthetase;

wherein each of said introduced DNAs is operatively linked to
15 regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of
propionyl-CoA compared to that in cells of a corresponding, wild-type plant
not comprising said introduced DNAs.

20

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

25

homoserine dehydrogenase;

threonine synthase;

30

a wild-type or deregulated threonine deaminase;

pyruvate oxidase; and

an acyl-CoA synthetase;

- 5 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- wherein cells of said plant produce an increased amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant
10 not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

- 15 a wild-type or deregulated threonine deaminase; and

- an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate
20 dehydrogenase complex in said plant;

 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- 25 wherein cells of said plant produce an increased amount of propionyl-Co A compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

- A plant, the genome of which comprises introduced DNAs encoding
30 the following enzymes:

threonine synthase;

a wild-type or deregulated threonine deaminase; and

5 an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

10 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of propionyl-Co A compared to that in cells of a corresponding, wild-type plant

15 not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

20 a wild-type or deregulated aspartate kinase;

a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved
25 binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

wherein each of said introduced DNAs is operatively linked to
30 regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

5 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

10 threonine synthase;

a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved
15 binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

wherein each of said introduced DNAs is operatively linked to
20 regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

25

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

30

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;
threonine synthase;

a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of
5 propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

10

a wild-type or deregulated threonine deaminase;

pyruvate oxidase; and

15

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

20

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant
25 not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

30

threonine synthase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase; and

5 an α -ketoacid decarboxylase E1 subunit that exhibits improved
binding and decarboxylating activities with α -ketobutyrate compared with
pyruvate decarboxylase E1 subunit naturally present in pyruvate
dehydrogenase complex in said plant;

10 wherein each of said introduced DNAs is operatively linked to
regulatory signals that cause expression of said introduced DNAs; and

 wherein cells of said plant produce an increased amount of
propionyl-CoA compared to that in cells of a corresponding, wild-type plant
15 not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding
the following enzymes:

20 a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

an acyl-CoA synthetase; and

25

 an α -ketoacid decarboxylase E1 subunit that exhibits improved
binding and decarboxylating activities with α -ketobutyrate compared with
pyruvate decarboxylase E1 subunit naturally present in pyruvate
dehydrogenase complex in said plant;

30

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of
5 propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

10

threonine synthase;

a wild-type or deregulated threonine deaminase;

15

pyruvate oxidase;

an acyl-CoA synthetase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved
20 binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

wherein each of said introduced DNAs is operatively linked to
25 regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

30

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

5

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

10

pyruvate oxidase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

15

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

20

wherein cells of said plant produce an increased amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

25

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

30

threonine synthase;

[illegible][illegible][illegible][illegible][illegible][illegible][illegible]

F **B** **E** **L** **I** **N** **G** **A** **T** **I** **O** **N** **S**

[illegible]

F **B** **E** **L** **I** **N** **G** **A** **S** **T** **R** **O** **N** **G** **M** **A** **N** **D** **A** **T** **E** **S**

[illegible]

pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

wherein each of said introduced DNAs is operatively linked to
5 regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

10

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated threonine deaminase;

15

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

20

threonine synthase;

pyruvate oxidase;

an acyl-CoA synthetase; and

25

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

30

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce an increased amount of
5 propionyl-CoA compared to that in cells of a corresponding, wild-type plant not comprising said introduced DNAs.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

10

a wild-type or deregulated aspartate kinase;

a wild-type or deregulated threonine deaminase;

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
20 to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxy-
25 valeryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

5

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

10

threonine synthase;

a wild-type or deregulated threonine deaminase;

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

25

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

5

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

10

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

20

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

25

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

5

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

10

homoserine dehydrogenase;

threonine synthase;

15

a wild-type or deregulated threonine deaminase;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

20

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

25

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating

β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to
5 regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding
10 the following enzymes:

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

20 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA
25 and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating

β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to
5 regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding
10 the following enzymes:

threonine synthase;

a wild-type or deregulated threonine deaminase;

15

pyruvate oxidase;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing
20 acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and
propionyl-CoA to produce β -ketovaleryl-CoA;

25

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA
and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -
hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

- 5 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

- 10 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated threonine deaminase;

- 15 pyruvate oxidase;

an acyl-CoA synthetase;

- 20 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 25 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

FILED FOR RECORD

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

5 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

10 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

threonine synthase;

15 a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

an acyl-CoA synthetase;

20

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

25 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 5 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

10 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

15 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

20 homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

- 25 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

5 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating
10 β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

15 wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

20 a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

25 threonine synthase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

30

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- 10 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 15 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- 20 wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

- 25 a wild-type or deregulated aspartate kinase;
- homoserine dehydrogenase;
- a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

an acyl-CoA synthetase;

- 5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 10 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- 15 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 20 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

25 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

threonine synthase;

5

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

10

an acyl-CoA synthetase;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

20

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating

25

β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

5

a wild-type or deregulated threonine deaminase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with
10 pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing
15 acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

20

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

25

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

10

[illegible]

15

20

25

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

- 5 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

- 10 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

- 15 a wild-type or deregulated threonine deaminase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate

- 20 dehydrogenase complex in said plant;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 25 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 5 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to
10 regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding
15 the following enzymes:

a wild-type or deregulated aspartate kinase;

threonine synthase;

20

a wild-type or deregulated threonine deaminase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved
binding and decarboxylating activities with α -ketobutyrate compared with
25 pyruvate decarboxylase E1 subunit naturally present in pyruvate
dehydrogenase complex in said plant;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- 10 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 15 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- 20 wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

- 25 a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

5

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

10

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

15
20

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

25

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

5 threonine synthase;

a wild-type or deregulated threonine deaminase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved
10 binding and decarboxylating activities with α -ketobutyrate compared with
pyruvate decarboxylase E1 subunit naturally present in pyruvate
dehydrogenase complex in said plant;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
15 to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing
acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and
20 propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA
and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxy-
valeryl-CoA, respectively; and

25 a polyhydroxyalkanoate synthase capable of incorporating
 β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV)
copolymer;

wherein each said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

5

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated threonine deaminase;

10

pyruvate oxidase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

20

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

25

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

5 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

10 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

threonine synthase;

15 a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved
20 binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
25 to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

0177 0178 0179 0180 0181 0182 0183 0184 0185 0186 0187 0188 0189 0190 0191 0192 0193 0194 0195 0196 0197 0198 0199 0200 0201 0202 0203 0204 0205 0206 0207 0208 0209 0210 0211 0212 0213 0214 0215 0216 0217 0218 0219 0220 0221 0222 0223 0224 0225 0226 0227 0228 0229 0230 0231 0232 0233 0234 0235 0236 0237 0238 0239 0240 0241 0242 0243 0244 0245 0246 0247 0248 0249 0250 0251 0252 0253 0254 0255 0256 0257 0258 0259 0260 0261 0262 0263 0264 0265 0266 0267 0268 0269 0270 0271 0272 0273 0274 0275 0276 0277 0278 0279 0280 0281 0282 0283 0284 0285 0286 0287 0288 0289 0290 0291 0292 0293 0294 0295 0296 0297 0298 0299 0300 0301 0302 0303 0304 0305 0306 0307 0308 0309 0310 0311 0312 0313 0314 0315 0316 0317 0318 0319 0320 0321 0322 0323 0324 0325 0326 0327 0328 0329 0330 0331 0332 0333 0334 0335 0336 0337 0338 0339 0340 0341 0342 0343 0344 0345 0346 0347 0348 0349 0350 0351 0352 0353 0354 0355 0356 0357 0358 0359 0360 0361 0362 0363 0364 0365 0366 0367 0368 0369 0370 0371 0372 0373 0374 0375 0376 0377 0378 0379 0380 0381 0382 0383 0384 0385 0386 0387 0388 0389 0390 0391 0392 0393 0394 0395 0396 0397 0398 0399 0400 0401 0402 0403 0404 0405 0406 0407 0408 0409 0410 0411 0412 0413 0414 0415 0416 0417 0418 0419 0420 0421 0422 0423 0424 0425 0426 0427 0428 0429 0430 0431 0432 0433 0434 0435 0436 0437 0438 0439 0440 0441 0442 0443 0444 0445 0446 0447 0448 0449 0450 0451 0452 0453 0454 0455 0456 0457 0458 0459 0460 0461 0462 0463 0464 0465 0466 0467 0468 0469 0470 0471 0472 0473 0474 0475 0476 0477 0478 0479 0480 0481 0482 0483 0484 0485 0486 0487 0488 0489 0490 0491 0492 0493 0494 0495 0496 0497 0498 0499 0500 0501 0502 0503 0504 0505 0506 0507 0508 0509 0510 0511 0512 0513 0514 0515 0516 0517 0518 0519 0520 0521 0522 0523 0524 0525 0526 0527 0528 0529 0530 0531 0532 0533 0534 0535 0536 0537 0538 0539 0540 0541 0542 0543 0544 0545 0546 0547 0548 0549 0550 0551 0552 0553 0554 0555 0556 0557 0558 0559 0560 0561 0562 0563 0564 0565 0566 0567 0568 0569 0570 0571 0572 0573 0574 0575 0576 0577 0578 0579 0580 0581 0582 0583 0584 0585 0586 0587 0588 0589 0590 0591 0592 0593 0594 0595 0596 0597 0598 0599 0600 0601 0602 0603 0604 0605 0606 0607 0608 0609 0610 0611 0612 0613 0614 0615 0616 0617 0618 0619 0620 0621 0622 0623 0624 0625 0626 0627 0628 0629 0630 0631 0632 0633 0634 0635 0636 0637 0638 0639 0640 0641 0642 0643 0644 0645 0646 0647 0648 0649 0650 0651 0652 0653 0654 0655 0656 0657 0658 0659 0660 0661 0662 0663 0664 0665 0666 0667 0668 0669 0670 0671 0672 0673 0674 0675 0676 0677 0678 0679 0680 0681 0682 0683 0684 0685 0686 0687 0688 0689 0690 0691 0692 0693 0694 0695 0696 0697 0698 0699 0700 0701 0702 0703 0704 0705 0706 0707 0708 0709 0710 0711 0712 0713 0714 0715 0716 0717 0718 0719 0720 0721 0722 0723 0724 0725 0726 0727 0728 0729 0730 0731 0732 0733 0734 0735 0736 0737 0738 0739 0740 0741 0742 0743 0744 0745 0746 0747 0748 0749 0750 0751 0752 0753 0754 0755 0756 0757 0758 0759 0760 0761 0762 0763 0764 0765 0766 0767 0768 0769 0770 0771 0772 0773 0774 0775 0776 0777 0778 0779 0780 0781 0782 0783 0784 0785 0786 0787 0788 0789 0790 0791 0792 0793 0794 0795 0796 0797 0798 0799 0800 0801 0802 0803 0804 0805 0806 0807 0808 0809 0810 0811 0812 0813 0814 0815 0816 0817 0818 0819 0820 0821 0822 0823 0824 0825 0826 0827 0828 0829 0830 0831 0832 0833 0834 0835 0836 0837 0838 0839 0840 0841 0842 0843 0844 0845 0846 0847 0848 0849 0850 0851 0852 0853 0854 0855 0856 0857 0858 0859 0860 0861 0862 0863 0864 0865 0866 0867 0868 0869 0870 0871 0872 0873 0874 0875 0876 0877 0878 0879 0880 0881 0882 0883 0884 0885 0886 0887 0888 0889 0890 0891 0892 0893 0894 0895 0896 0897 0898 0899 0900 0901 0902 0903 0904 0905 0906 0907 0908 0909 0910 0911 0912 0913 0914 0915 0916 0917 0918 0919 0920 0921 0922 0923 0924 0925 0926 0927 0928 0929 0930 0931 0932 0933 0934 0935 0936 0937 0938 0939 0940 0941 0942 0943 0944 0945 0946 0947 0948 0949 0950 0951 0952 0953 0954 0955 0956 0957 0958 0959 0960 0961 0962 0963 0964 0965 0966 0967 0968 0969 0970 0971 0972 0973 0974 0975 0976 0977 0978 0979 0980 0981 0982 0983 0984 0985 0986 0987 0988 0989 0990 0991 0992 0993 0994 0

1.

10

15

20

25

• • •

pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
5 to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing
acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and
10 propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA
and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxy-
valeryl-CoA, respectively; and
15

a polyhydroxyalkanoate synthase capable of incorporating
 β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV)
copolymer;

20 wherein each of said introduced DNAs is operatively linked to
regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

25 A plant, the genome of which comprises introduced DNAs encoding
the following enzymes:

threonine synthase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

5 an acyl-CoA synthetase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved
binding and decarboxylating activities with α -ketobutyrate compared with
pyruvate decarboxylase E1 subunit naturally present in pyruvate
10 dehydrogenase complex in said plant;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing
acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or
15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and
propionyl-CoA to produce β -ketovaleryl-CoA;

20 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA
and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxy-
valeryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating
25 β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV)
copolymer;

TOOBER 13618A

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

5

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

10

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

15

pyruvate oxidase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

20

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

25

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

5 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

10 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

15 A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

20

threonine synthase;

a wild-type or deregulated threonine deaminase;

25

pyruvate oxidase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate
30 dehydrogenase complex in said plant;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- 10 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 15 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- 20 wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

- 25 a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

an acyl-CoA synthetase;

5 an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

10 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
15 to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxy-
20 valeryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

25

wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

A plant, the genome of which comprises introduced DNAs encoding the following enzymes:

5

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

10

threonine synthase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

15

an acyl-CoA synthetase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said plant;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

5 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

10 wherein each of said introduced DNAs is operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein cells of said plant produce P(3HB-co-3HV) copolymer.

15 Any of the foregoing plants, wherein each of said introduced DNAs is further operatively linked to a plastid transit peptide coding region capable of directing transport of said enzyme encoded thereby into a plastid.

20 A method of producing P(3HB-co-3HV)copolymer, comprising growing any of the foregoing plants, the genome of which comprises introduced DNAs encoding a β -ketothiolase, a β -ketoacyl-CoA reductase, and a polyhydroxyalkanoate synthase, and recovering said P(3HB-co-3HV) copolymer produced thereby.

25 The foregoing method, wherein said P(3HB-co-3HV)copolymer is recovered from leaves or seeds of said plant.

A plant cell containing P(3HB-co-3HV) copolymer.

A plant comprising cells containing P(3HB-co-3HV) copolymer.

Seeds of the foregoing plant.

The foregoing plant, wherein said cells are located in leaves or seeds of said plant.

5

A plant, the genome of which comprises introduced DNAs encoding the enzymes PhbA, PhbB, and PhbC;

wherein each of said introduced DNAs is operatively linked to a
10 plastid transit peptide coding region capable of directing transport of said enzymes into a plastid, and regulatory signals that cause expression of said introduced DNAs in seeds of said plant; and

15

wherein P(3HB) homopolymer is produced in seeds of said plant.

A method of producing P(3HB) homopolymer, comprising growing the foregoing plant, and recovering said P(3HB) homopolymer produced thereby.

20

The foregoing method, wherein said P(3HB) homopolymer is recovered from seeds of said plant.

25

A bacterium, the genome of which comprises introduced DNA encoding a wild-type or deregulated aspartate kinase;

wherein said introduced DNA is operatively linked to regulatory signals that cause expression of said introduced DNA; and

wherein said bacterium produces an elevated amount of threonine
30 compared to that in a corresponding wild-type bacterium not comprising said introduced DNA.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase; and

5

threonine synthase;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

10

wherein said bacterium produces an elevated amount of threonine compared to that in a corresponding wild-type bacterium not comprising said introduced DNAs.

15

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase; and

20

homoserine dehydrogenase;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

25

wherein said bacterium produces an elevated amount of threonine compared to that in a corresponding wild-type bacterium not comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

30

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase; and

5 threonine synthase;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

10 wherein said bacterium produces an elevated amount of threonine compared to that in a corresponding wild-type bacterium not comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs
15 encoding the following enzymes:

a wild-type or deregulated aspartate kinase; and

a wild-type or deregulated threonine deaminase;

20 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an elevated amount of α -keto-
25 butyrate compared to that in a corresponding wild-type bacterium not comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

30 a wild-type or deregulated aspartate kinase;

threonine synthase; and

a wild-type or deregulated threonine deaminase;

- 5 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

 wherein said bacterium produces an elevated amount of α -keto-butyrate compared to that in a corresponding wild-type bacterium not
10 comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

- 15 a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase; and

a wild-type or deregulated threonine deaminase;

20

 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

 wherein said bacterium produces an increased amount of α -keto-butyrate compared to that in a wild-type bacterium not comprising said
25 introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

30

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

threonine synthase; and

5 a wild-type or deregulated threonine deaminase;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

10 wherein said bacterium produces an increased amount of α -keto-butyrate compared to that in a wild-type bacterium not comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs
15 encoding the following enzymes:

a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved
20 binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex of said bacterium;

wherein each of said introduced DNAs is operatively linked to
25 regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an elevated amount of propionyl-CoA compared to that in a corresponding wild-type bacterium not comprising said introduced DNAs.

The foregoing bacterium, wherein said α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate is a branched chain α -ketoacid decarboxylase E1 subunit such as that from bovine kidney, *Pseudomonas putida*, or *Bacillus subtilis*.

5

A bacterium, the genome of which comprises introduced DNAs encoding:

a wild-type or deregulated threonine deaminase;

10

a pyruvate decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium; and

15

a dihydrolipoyl transacylase E2 subunit exhibiting improved transacylase activity with α -ketobutyrate compared with dihydrolipoyl transacetylase E2 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

20

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an elevated amount of propionyl-CoA compared to that in a corresponding wild-type bacterium not comprising said introduced DNAs.

25

The foregoing bacterium, wherein said α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate is a branched chain α -ketoacid decarboxylase E1 subunit

30

such as that from bovine kidney, *Pseudomonas putida*, or *Bacillus subtilis*, and said dihydrolipoyl transacylase E2 subunit exhibiting improved transacylase activity with α -ketobutyrate is one such as that from bovine kidney, *Pseudomonas putida*, or *Bacillus subtilis*.

5

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

10

a wild-type or deregulated threonine deaminase; and
pyruvate oxidase;

15

wherein said introduced DNAs are operably linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an elevated amount of propionyl-CoA compared that in a corresponding wild-type bacterium not comprising said introduced DNAs.

20

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

threonine synthase;

25

a wild-type or deregulated threonine deaminase; and
pyruvate oxidase;

wherein said introduced DNAs are operably linked to regulatory
signals that cause expression of said introduced DNAs; and

F032630162460

wherein said bacterium produces an elevated amount of propionyl-CoA compared that in a corresponding wild-type bacterium not comprising said introduced DNAs.

- 5 The foregoing bacterium, wherein said pyruvate oxidase is encoded by *E. coli poxB*.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

10

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

25

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

5

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

threonine synthase;

10

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxy-valeryl-CoA, respectively; and

25

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

5

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated threonine deaminase;

10

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

15

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

25

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

5

threonine synthase;

a wild-type or deregulated threonine deaminase;

10

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

15 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

20 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

25

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

Approved for Release

The foregoing bacterium, wherein said β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA is BktB.

- 5 The foregoing bacterium, wherein said β -ketoacyl-CoA reductase is acetoacetyl-CoA reductase.

- The foregoing bacterium, wherein said polyhydroxyalkanoate synthase is obtainable from a microorganism selected from the group
- 10 consisting of *Alcaligenes eutrophus*, *Alcaligenes faecalis*, *Aphanotheca* sp., *Azotobacter vinelandii*, *Bacillus cereus*, *Bacillus megaterium*, *Beijerinckia indica*, *Derxia gummosa*, *Methylobacterium* sp., *Microcoleus* sp., *Nocardia corallina*, *Pseudomonas cepacia*, *Pseudomonas extorquens*, *Pseudomonas oleovorans*, *Rhodobacter sphaeroides*, *Rhodobacter capsulatus*,
- 15 *Rhodospirillum rubrum*, and *Thiocapsa pfennigii*.

A bacterium, the genome of which comprises introduced DNA encoding a wild-type or deregulated threonine deaminase;

- 20 wherein said introduced DNA is operatively linked to regulatory signals that cause expression of said introduced DNA; and

- wherein said bacterium produces an increased amount of α -keto-butyrate compared to that in a corresponding wild-type bacterium not
- 25 comprising said introduced DNA.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

- 30 threonine synthase; and

a wild-type or deregulated threonine deaminase;

wherein said introduced DNAs are operatively linked to regulatory
5 signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of α -keto-
butyrate compared to that in a corresponding wild-type bacterium not
comprising said introduced DNAs.

10

A bacterium, the genome of which comprises introduced DNAs
encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

15

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase; and

20

pyruvate oxidase;

wherein said introduced DNAs are operatively linked to regulatory
signals that cause expression of said introduced DNAs; and

25 wherein said bacterium produces an increased amount of propionyl-
CoA compared to that in a corresponding, wild-type bacterium not
comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs
30 encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

threonine synthase;

5

a wild-type or deregulated threonine deaminase; and
pyruvate oxidase;

wherein said introduced DNAs are operatively linked to regulatory
10 signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-
CoA compared to that in a corresponding, wild-type bacterium not
comprising said introduced DNAs.

15

A bacterium, the genome of which comprises introduced DNAs
encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

20

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

25

pyruvate oxidase; and

an acyl-CoA synthetase;

wherein said introduced DNAs are operatively linked to regulatory
30 signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

5 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

10 homoserine dehydrogenase;

threonine synthase;

a wild-type or deregulated threonine deaminase;

15 pyruvate oxidase; and

an acyl-CoA synthetase;

20 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not
25 comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

30 a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

5

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-Co A compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

15

threonine synthase;

a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-Co A compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

5

a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

10

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

15

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

20 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

25

threonine synthase;

a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with

30

pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

wherein said introduced DNAs are operatively linked to regulatory
5 signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

10

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

15

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase; and

20

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

25

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not

30 comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

5 a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

threonine synthase;

10 a wild-type or deregulated threonine deaminase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate

15 dehydrogenase complex in said bacterium;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

20 wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated threonine deaminase;

pyruvate oxidase; and

30

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

5

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

15

threonine synthase;

a wild-type or deregulated threonine deaminase;

20

pyruvate oxidase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

25

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

T03290-4032400

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

5 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated threonine deaminase;

10 pyruvate oxidase;

an acyl-CoA synthetase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved
15 binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

wherein said introduced DNAs are operatively linked to regulatory
20 signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

25

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

threonine synthase;

30

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

an acyl-CoA synthetase; and

5 an α -ketoacid decarboxylase E1 subunit that exhibits improved
binding and decarboxylating activities with α -ketobutyrate compared with
pyruvate decarboxylase E1 subunit naturally present in pyruvate
dehydrogenase complex in said bacterium;

10 wherein said introduced DNAs are operatively linked to regulatory
signals that cause expression of said introduced DNAs; and

 wherein said bacterium produces an increased amount of propionyl-
CoA compared to that in a corresponding, wild-type bacterium not
15 comprising said introduced DNAs.

A bacterium, the genome of which comprises introduced DNAs
encoding the following enzymes:

20 a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

25 pyruvate oxidase; and

 an α -ketoacid decarboxylase E1 subunit that exhibits improved
binding and decarboxylating activities with α -ketobutyrate compared with

437000 1000000

pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

wherein said introduced DNAs are operatively linked to regulatory
5 signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

10

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

15

homoserine dehydrogenase;

threonine synthase;

20

a wild-type or deregulated threonine deaminase;

pyruvate oxidase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved
25 binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

wherein said introduced DNAs are operatively linked to regulatory
30 signals that cause expression of said introduced DNAs; and

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

5 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

10 homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

15 an acyl-CoA synthetase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with
20 pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and
25

wherein said bacterium produces an increased amount of propionyl-CoA compared to that in a corresponding, wild-type bacterium not comprising said introduced DNAs.

30 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

5 threonine synthase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

10

an acyl-CoA synthetase; and

an α -ketoacid decarboxylase E1 subunit that exhibits improved
binding and decarboxylating activities with α -ketobutyrate compared with
15 pyruvate decarboxylase E1 subunit naturally present in pyruvate
dehydrogenase complex in said bacterium;

wherein said introduced DNAs are operatively linked to regulatory
signals that cause expression of said introduced DNAs; and

20

wherein said bacterium produces an increased amount of propionyl-
CoA compared to that in a corresponding, wild-type bacterium not
comprising said introduced DNAs.

25 A bacterium, the genome of which comprises introduced DNAs
encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

30

a wild-type or deregulated threonine deaminase;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA
10 and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- a polyhydroxyalkanoate synthase capable of incorporating
 β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV)
15 copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- 20 wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

- 25 a wild-type or deregulated aspartate kinase;

threonine synthase;

a wild-type or deregulated threonine deaminase;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA
10 and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV)
15 copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- 20 wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

- 25 a wild-type, partially, or totally lysine feedback inhibition insensitive aspartate kinase;

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing

5 acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

10

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

15

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

20 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

25 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type, partially, or totally lysine feedback inhibition insensitive aspartate kinase;

homoserine dehydrogenase;

threonine synthase;

5 a wild-type or deregulated threonine deaminase;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

10

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

15 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating
20 β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

25

wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

13618-21-38

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

10 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

15 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

20 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

25 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

threonine synthase;

FBI/DOJ - 100-443889

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

10 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

15 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

20 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

25 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

an acyl-CoA synthetase;

- 5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 10 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- 15 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 20 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

25 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

threonine synthase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

5 an acyl-CoA synthetase;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

10

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

15 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating
20 β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

25

wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

rule

CoA

rule
acet

acc
A an

corpo
p(3E

ake
and

7) c

acet
prop
iol
cet
pro
cyl

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

- 5 a wild-type or deregulated aspartate kinase;
- homoserine dehydrogenase;
- threonine synthase;
- 10 a wild-type or deregulated threonine deaminase;
- pyruvate oxidase;
- a β -ketothiolase capable of condensing two molecules of acetyl-CoA
- 15 to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or
- a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and
- 20 propionyl-CoA to produce β -ketovaleryl-CoA;
- a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and
- 25 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

5

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

10

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

15

pyruvate oxidase;

an acyl-CoA synthetase;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA

20 to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and
25 propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

5 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

10 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

15 homoserine dehydrogenase;

threonine synthase;

a wild-type or deregulated threonine deaminase;

20

pyruvate oxidase;

an acyl-CoA synthetase;

25 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- 5 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 10 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- 15 wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

- 20 a wild-type or deregulated threonine deaminase;

- an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;
- 25

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- 10 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 15 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- 20 wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

- 25 threonine synthase;

a wild-type or deregulated threonine deaminase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

5

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

10

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA
15 and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating
 β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV)
20 copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

25

wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

a wild-type or deregulated threonine deaminase;

5 an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

10 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
15 to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxy-
20 valeryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

25 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

5

a wild-type or deregulated aspartate kinase;

threonine synthase;

10

a wild-type or deregulated threonine deaminase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate

15 dehydrogenase complex in said bacterium;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

20

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

25

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating

Pub No. 13618-21

β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory
5 signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs
10 encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

15

a wild-type or deregulated threonine deaminase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved
binding and decarboxylating activities with α -ketobutyrate compared with
20 pyruvate decarboxylase E1 subunit naturally present in pyruvate
dehydrogenase complex in said bacterium;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing
25 acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and
propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

5 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

10 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

15 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

20

threonine synthase;

a wild-type or deregulated threonine deaminase;

25 an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- 10 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 15 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- 20 wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

- 25 a wild-type or deregulated threonine deaminase;
- pyruvate oxidase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

5

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

10

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

15

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

20

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

25

wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

FOUO "FOUO" FOUO

threonine synthase;

a wild-type or deregulated threonine deaminase;

5 pyruvate oxidase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate
10 dehydrogenase complex in said bacterium;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

15 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

20 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating
25 β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

5

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated threonine deaminase;

10

pyruvate oxidase;

an acyl-CoA synthetase;

15

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

20

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

25

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

5 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

10 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs
15 encoding the following enzymes:

threonine synthase;

a wild-type or deregulated threonine deaminase;

20

pyruvate oxidase;

an acyl-CoA synthetase;

25 an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- 5 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- 10 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 15 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

- 20 wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

- 25 a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with

- 5 pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

- a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing
10 acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

- a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

- 15 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 20 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

- wherein said introduced DNAs are operatively linked to regulatory
25 signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

- 5 a wild-type or deregulated aspartate kinase;
- homoserine dehydrogenase;
- threonine synthase;
- 10 a wild-type or deregulated threonine deaminase;
- pyruvate oxidase;
- an α -ketoacid decarboxylase E1 subunit that exhibits improved
- 15 binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;
- a β -ketothiolase capable of condensing two molecules of acetyl-CoA
- 20 to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or
- a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and
- 25 propionyl-CoA to produce β -ketovaleryl-CoA;
- a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

5 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

10 A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

a wild-type or deregulated aspartate kinase;

15 homoserine dehydrogenase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

20

an acyl-CoA synthetase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with
25 pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing
30 acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

5 a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

10 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

15 wherein said bacterium produces P(3HB-co-3HV) copolymer.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

20 a wild-type or deregulated aspartate kinase;

homoserine dehydrogenase;

25 threonine synthase;

a wild-type or deregulated threonine deaminase;

pyruvate oxidase;

30

an acyl-CoA synthetase;

an α -ketoacid decarboxylase E1 subunit that exhibits improved binding and decarboxylating activities with α -ketobutyrate compared with pyruvate decarboxylase E1 subunit naturally present in pyruvate dehydrogenase complex in said bacterium;

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

A method of producing P(3HB-co-3HV) copolymer, comprising growing any of the foregoing bacteria, the genome of which comprises introduced DNAs encoding a β -ketothiolase, a β -ketoacyl-CoA reductase, and a polyhydroxy-alkanoate synthase, and recovering said P(3HB-co-
5 3HV)copolymer produced thereby.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

10 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

a β -ketothiolase capable of condensing two molecules of acetyl-CoA
15 to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxy-
20 valeryl-CoA, respectively; and

a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

25

wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer.

The foregoing bacterium can be one that overproduces threonine. The genome thereof can further comprise, in addition to introduced DNAs
5 encoding the enzymes listed above, introduced DNA encoding a wild-type or deregulated threonine deaminase. Alternatively, the foregoing bacterium can be one that produces propionate or propionyl-CoA at levels useful for producing P(3HB-co-3HV) copolymer.

10 A method of producing P(3HB-co-3HV) copolymer, comprising:

culturing any of the four foregoing mentioned bacteria under conditions and for a time conducive to the formation of P(3HB-co-3HV) copolymer; and

15 recovering said P(3HB-co-3HV) copolymer produced thereby.

A bacterium, the genome of which comprises introduced DNAs encoding the following enzymes:

20 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA and a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA; or

25 a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, and capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA;

a β -ketoacyl-CoA reductase capable of reducing acetoacetyl-CoA and β -ketovaleryl-CoA to produce β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA, respectively; and

- 5 a polyhydroxyalkanoate synthase capable of incorporating β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV) copolymer;

10 wherein said introduced DNAs are operatively linked to regulatory signals that cause expression of said introduced DNAs; and

wherein said bacterium produces P(3HB-co-3HV) copolymer when grown on a medium comprising propionic acid.

- 15 A method of producing P(3HB-co-3HV) copolymer, comprising:

culturing the foregoing bacterium on a medium comprising propionic acid under conditions and for a time conducive to the formation of P(3HB-co-3HV) copolymer, and

20

recovering said P(3HB-co-3HV) copolymer produced thereby.

- 25 A bacterial cell containing P(3HB-co-3HV) copolymer, wherein said P(3HB-co-3HV) copolymer is produced within said bacterial cell as a result of expression of at least one DNA sequence introduced therein that encodes an enzyme necessary for P(3HB-co-3HV) copolymer synthesis.

- 30 The foregoing bacterial cell, wherein said at least one DNA sequence is selected from the group consisting of a DNA sequence encoding a β -keto thiolase capable of condensing two molecules of acetyl-CoA to produce

acetoacetyl-Co-A, a DNA sequence encoding a β -ketothiolase capable of
condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-CoA, a
DNA sequence encoding a β -ketoacyl-CoA reductase, a DNA sequence
encoding a polyhydroxyalkanoate synthase capable of incorporating
5 β -hydroxybutyryl-CoA and β -hydroxyvaleryl-CoA into P(3HB-co-3HV)
copolymer, and combinations thereof.

The foregoing bacterial cell, further comprising a DNA sequence
introduced therein that encodes a wild-type or deregulated threonine
10 deaminase, wherein said DNA sequence encoding said wild-type or
deregulated threonine deaminase is expressed.

A method for transforming canola, comprising:

15 (a) preparing a stem explant from a canola plant by:

(i) removing leaves and buds along the stem and removing 4-5
inches of said stem below the flower buds; and

20 (ii) cutting said 4-5 inches of stem into segments;

(b) inserting DNA to be introduced into said explant of step (a) by
inoculating said explant with a disarmed *Agrobacterium tumefaciens* vector
containing said DNA;

25

(c) culturing said explant of step (b) in the basal-side down
orientation;

(d) selecting transformed explant tissue; and

30

(e) regenerating a differentiated transformed plant from said transformed explant tissue of step (d).

A method for transforming soybean, comprising:

5

(a) preparing a cotyledon explant from a soybean seedling by:

(i) incubating said seedling at about 0°C to about 10°C for at least 24 hours;

10

(ii) removing the hypocotyl region by cutting in the region of from about 0.2 to about 1.5 cm below the cotyledonary node;

(ii) splitting and completely separating the remaining
15 attached hypocotyl segment, also thereby separating the two cotyledons;

(iii) removing the epicotyl from the cotyledon to which it remains attached; and

20

(iv) wounding the cotyledon in the region of said axillary bud;

(b) inserting DNA to be introduced into said explant of step (a) by inoculating at least the region adjacent to the axillary bud of the explant with a disarmed *Agrobacterium tumefaciens* vector containing said DNA;

25

(c) selecting transformed explant tissue; and

(d) regenerating a differentiated transformed plant from said transformed explant tissue of step (c).

30

Approved for Release by NSA on 09-11-2013 pursuant to E.O. 13526

A soybean explant prepared by steps (a)(i)-(a)(iv) of the preceding method.

Soybean tissue prepared from a seedling cotyledon pair containing
5 an epicotyl, axillary buds, and hypocotyl tissue, comprising a single
cotyledon containing an axillary bud and associated hypocotyl segment
extending from about 0.2 to about 1.5 cm below the cotyledonary node;

wherein said associated hypocotyl segment is completely separated
10 from its adjacent hypocotyl segment attached to the remaining cotyledon,
thus separating said cotyledons;

wherein said epicotyl has been removed from the cotyledon to which
it is attached;

15 wherein the cotyledon is wounded in the region of said axillary bud;
and

wherein said seedling has been incubated at a temperature of from
20 about 0°C to about 10°C for at least about 24 hours prior to preparing said
soybean tissue.

Further scope of the applicability of the present invention will
become apparent from the detailed description and drawings provided
25 below. However, it should be understood that the detailed description and
specific examples, while indicating preferred embodiments of the present
invention, are given by way of illustration only since various changes and
modifications within the spirit and scope of the invention will become
apparent to those skilled in the art from this detailed description.

Brief Description of the Drawings

The above and other aspects, features, and advantages of the present invention will be better understood from the following detailed description taken in conjunction with the accompanying drawings, all of which are given by way of illustration only and are not limitative of the present invention.

For the sake of brevity, a description of many of the common genetic elements present in each of the plasmid figures is not included in each figure legend. These genetic elements include Amp (ampicillin bacterial resistance); Spec or Spc/Str (spectinomycin bacterial resistance); Kan or Kan903 (kanamycin bacterial resistance); P-Sp6 (Sp6 RNA polymerase promoter); P-T7 (T7 RNA polymerase promoter); ori-pUC; ori-327; and ori-322 (each designation refers to the ColE1 plasmid origin of DNA replication of *E. coli*); ori-M13 (M13 phage origin of replication); ori-pACYC (plasmid origin of DNA replication of *E. coli*); ptrc and ptac (IPTG-inducible *E. coli* Trp/Lac fusion promoters); G10 (T7 gene 10 translational enhancing leader); preA (*recA* SOS responsive promoter); e35S, p7S, and P-FMV (promoters for plant expression); ARABSSU1A or PEPSPS-transit:2 (chloroplast transit peptides); E9 3' and NOS 3' (plant polyadenylation/transcription termination signals); ori-V (*Agrobacterium* origin of replication).

Figure 1 shows the biochemical steps involved in the production of PHB from acetyl-CoA catalyzed by the *A. eutrophus* PHB biosynthetic enzymes.

Figure 2 shows the biochemical steps involved in the production of P(3HB-co-3HV) copolymer from acetyl-CoA and propionyl-CoA catalyzed by PHA biosynthetic enzymes of *A. eutrophus*.

Figure 3 summarizes the pathways discussed herein that are involved in the production of P(3HB-co-3HV) copolymer.

Figure 4 shows the structure of pMON25659. pMON25659
5 contains the wild-type *E. coli* biosynthetic threonine deaminase (*ilvA*) gene derived by PCR. The nucleotide sequence of the wild-type biosynthetic threonine deaminase gene is shown in SEQ ID NO: 1.

Figure 5 shows the structure of pMON25660. pMON25660 was
10 used for the overexpression of the wild-type biosynthetic threonine deaminase (IlvA) from the ptrc promoter in *E. coli*.

Figure 6 shows the structure of pMON25676. pMON25676
contains the same BamHI fragment as is present in pMON25659. This
15 plasmid was used for preparation of ssDNA for *in vitro* mutagenesis.

Figure 7 shows the structure of pMON25679. pMON25679
contains the *ilvA219* (L447F) gene encoding the mutant *E. coli*
biosynthetic threonine deaminase created by site-directed mutagenesis of
20 the wild-type threonine deaminase gene contained in plasmid pMON25676.
The nucleotide sequence of the *ilvA219* biosynthetic threonine deaminase gene is shown in SEQ ID NO:5.

Figure 8 shows the structure of pMON25680. pMON25680
25 contains the *ilvA466* (L481F) gene encoding the mutant *E. coli*
biosynthetic threonine deaminase created by site-directed mutagenesis of
the wild-type threonine deaminase gene contained in plasmid pMON25676.
The nucleotide sequence of the *ilvA466* biosynthetic threonine deaminase gene is shown in SEQ ID NO:7.

Figure 9 shows the structure of pMON25681. pMON25681 contains the *ilvA219/466* (L447F/L481F) gene encoding the mutant *E. coli* biosynthetic threonine deaminase created by site-directed mutagenesis of the wild-type threonine deaminase gene contained in plasmid pMON25676.

- 5 The nucleotide sequence of the *ilvA219/466* biosynthetic threonine deaminase gene is shown in SEQ ID NO: 8.

Figure 10 shows the structure of pMON25682. pMON25682 was used for the overexpression in *E. coli* of the *IlvA219* (L447F) mutant biosynthetic threonine deaminase from the *ptrc* promoter.

Figure 11 shows the structure of pMON25683. pMON25683 was used for the overexpression in *E. coli* of the *IlvA466* (L481F) mutant biosynthetic threonine deaminase from the *ptrc* promoter.

15 Figure 12 shows the structure of pMON25684. pMON25684 was used for the overexpression of the *IlvA219/466* (L447F/L481F) mutant biosynthetic threonine deaminase in *E. coli*.

20 Figure 13 shows the structure of pMON25663. pMON25663 was used for transient expression of the wild-type *E. coli* threonine deaminase (*IlvA*) in tobacco protoplasts.

Figure 14 shows the structure of pMON25686. pMON25686 was used for transient expression of the *IlvA219* (L447F) mutant threonine deaminase in tobacco protoplasts.

Figure 15 shows the structure of pMON25687. pMON25687 was used for transient expression of the *IlvA466* (L481F) mutant threonine deaminase in tobacco protoplasts.

For publication

Figure 16 shows the structure of pMON25688. pMON25688 was used for transient expression of the IlvA219/466 (L447F/L481F) mutant threonine deaminase in tobacco protoplasts.

5 Figure 17 shows the expression, confirmed by Western blot analysis, of wild-type and mutant biosynthetic *E. coli* threonine deaminases in tobacco protoplasts via electroporation with plasmids pMON26663, pMON25686, pMON25687, and pMON25688.

10 Immunoreactive bands corresponding to the predicted molecular weight of each threonine deaminase (56kDa) were observed in replicates, designated "1" and "2".

15 Figure 18 shows the structure of pMON25668. pMON25668 is the *Agrobacterium* plant transformation plasmid used to test expression of the wild-type *E. coli* threonine deaminase (IlvA) in soybean callus. The *phbA* gene on plasmid pMON25668 was not required for the expression of threonine deaminase.

20 Figure 19 shows the expression and activity of wild-type *E. coli* biosynthetic threonine deaminase in transformed soybean callus.

Figure 20 shows the structure of pMON25628. pMON25628 was used for the overexpression in *E. coli* of the *A. eutrophus* PhbB reductase from the *precA* promoter.

25

Figure 21 shows the structure of pMON25629. pMON25629 was used for the overexpression in *E. coli* of the *A. eutrophus* PhbC synthase from the *ptac* promoter.

Figure 22 shows the structure of pMON25636. pMON25636 was used for the overexpression in *E. coli* of the *A. eutrophus* PhbA β -keto-thiolase from the ptrc promoter.

5 Figure 23 shows the structure of pMON25626. pMON25626 was the *Agrobacterium* plant transformation vector containing the genetic elements for seed-specific, plastid-targeted expression of PhbB reductase and PhbC synthase used to transform canola and soybean.

10 Figure 24 shows the structure of pMON25638. pMON25638 was the *Agrobacterium* plant transformation vector containing the genetic elements for seed specific, plastid-targeted expression of PhbA β -keto-thiolase used to transform canola and soybean.

15 Figure 25 shows the structure of pMON25822. This plasmid is derived from the broad host-range vector pBBR1MCS-3 (Kovach, 1994), and expresses *A. eutrophus bktB* from the *A. eutrophus bktB* promoter. The *bktB* promoter is included on pMON25728, discussed in example 8.

20 Conventional methods of gene isolation, molecular cloning, vector construction, etc., are well known in the art and are summarized in Sambrook et al., 1989, and Ausubel et al., 1989. One skilled in the art can readily reproduce the plasmids vectors described above without undue experimentation employing these methods in conjunction with the cloning
25 information provided by the figures attached hereto. The various DNA sequences, fragments, etc., necessary for this purpose can be readily obtained as components of commercially available plasmids, or are otherwise well known in the art and publicly available.

The following detailed description of the invention is provided to aid those skilled in the art in practicing the present invention. Even so, the following detailed description should not be construed to unduly limit the present invention as modifications and variations in the embodiments
5 discussed herein can be made by those of ordinary skill in the art without departing from the spirit or scope of the present inventive discovery.

The references cited herein evidence the level of skill in the art to which the present invention pertains. The contents of each of these
10 references, including the references cited therein, are herein incorporated by reference in their entirety.

The following Examples describe a variety of different methods for enhancing the levels of threonine, α -ketobutyrate, and propionyl-CoA, the
15 latter being a direct precursor substrate in the synthesis of P(3HB-co-3HV) in bacteria and plants, and for producing PHA copolymers of differing C4/C5 compositions from various carbon sources.

Definitions

20 The following definitions are provided in order to aid those skilled in the art in understanding the detailed description of the present invention.

" β -ketoacyl-CoA reductase" refers to a β -ketoacyl-CoA reducing
25 enzyme that can convert a β -ketoacyl-CoA substrate to its corresponding β -hydroxyacyl-CoA product using, for example, NADH or NADPH as the reducing cosubstrate. An example is the PhbB acetoacetyl-CoA reductase of *A. eutrophus*.

“ β -ketothiolase” refers to an enzyme that catalyzes the thiolytic cleavage of a β -ketoacyl-CoA, requiring free CoA, to form two acyl-CoA molecules. However, the term “ β -ketothiolase” as used herein also refers to enzymes that catalyze the condensation of two acyl-CoA molecules to
5 form β -ketoacyl-CoA and free CoA, i.e., the reverse of the thiolytic cleavage reaction.

“CoA” refers to coenzyme A.

10 “C-terminal region” refers to the region of a peptide, polypeptide, or protein chain from the middle thereof to the end that carries the amino acid having a free α carboxyl group.

“Deregulated enzyme” refers to an enzyme that has been modified,
15 for example by mutagenesis, wherein the extent of feedback inhibition of the catalytic activity of the enzyme by a metabolite is reduced such that the enzyme exhibits enhanced activity in the presence of said metabolite compared to the unmodified enzyme. Some organisms possess deregulated forms of such enzymes as the naturally occurring, wild-type form.

20

The term “encoding DNA” refers to chromosomal DNA, plasmid DNA, cDNA, or synthetic DNA which codes on expression for any of the enzymes discussed herein.

25 The term “genome” as it applies to bacteria encompasses both the chromosome and plasmids within a bacterial host cell. Encoding DNAs of the present invention introduced into bacterial host cells can therefore be either chromosomally-integrated or plasmid-localized. The term “genome” as it applies to plant cells encompasses not only chromosomal DNA found
30 within the nucleus, but organelle DNA found within subcellular components

of the cell. DNAs of the present invention introduced into plant cells can therefore be either chromosomally-integrated or organelle-localized.

The terms "microbe" or "microorganism" refer to algae, bacteria,
5 fungi, and protozoa.

The term "mutein" refers to a mutant form of a peptide, polypeptide, or protein.

"N-terminal region" refers to the region of a peptide, polypeptide, or
10 protein chain from the amino acid having a free α amino group to the middle of the chain.

"Overexpression" refers to the expression of a polypeptide or protein encoded by a DNA introduced into a host cell, wherein said polypeptide or
15 protein is either not normally present in the host cell, or wherein said polypeptide or protein is present in said host cell at a higher level than that normally expressed from the endogenous gene encoding said polypeptide or protein.

20 The term "plastid" refers to the class of plant cell organelles that includes amyloplasts, chloroplasts, chromoplasts, elaioplasts, eoplasts, etioplasts, leucoplasts, and proplastids. These organelles are self-replicating, and contain what is commonly referred to as the "chloroplast genome," a circular DNA molecule that ranges in size from about 120 to
25 about 217 kb, depending upon the plant species, and which usually contains an inverted repeat region (Fosket, 1994).

The term "polyhydroxyalkanoate (PHA) synthase" refers to enzymes that convert β -hydroxyacyl-CoAs to polyhydroxyalkanoates and
30 free CoA.

The term “u” refers to an enzyme unit. One unit catalyzes the production of one μ mole of product per minute.

5

Example 1**Increased Production of Amino Acids in Bacterial and Plant Cells
Via Modification of the Activity of Aspartate Kinase, Homoserine
Dehydrogenase, and Threonine Synthase**

10 Biosynthesis of the aspartate family of amino acids in plants occurs in the plastids (Bryan, 1980). Figure 3 shows the enzymatic steps involved in the biosynthesis of P(3HB-co-3HV) copolymer from L-threonine via α -keto-butyrate. Both acetyl-CoA and propionyl-CoA are precursor substrates in the synthesis of P(3HB-co-3HV) as shown in this figure.

15

The involvement of L-threonine in the formation of α -ketobutyrate is described in detail below in Examples 2, 3, and 4. The present example describes methods for increasing the available pools of L-threonine in bacterial and plant cells for P(3HB-co-3HV) copolymer biosynthesis.

20

Overexpression of a threonine or lysine deregulated aspartate kinase (AK), the enzyme that catalyzes the first step in the biosynthesis of threonine, lysine, and methionine (Figure 3), results in an increase in the intracellular levels of free L-threonine in the leaf by 55% (Shaul and Galili, 25 1992a), and in the seed by 15-fold (Karchi et al., 1993).

A downstream enzyme in the aspartate pathway is dihydrodipicolinate synthase (DHPS), which catalyzes the first committed step in lysine synthesis (Figure 3). Plants that overexpress a deregulated 30 form of DHPS exhibit a 15- fold increase in free lysine (Shaul and Galili,

1992b). Plants expressing deregulated forms of both AK and DHPS exhibit a dramatic increase in free lysine, but lower levels of free threonine (Shaul and Galili, 1993). Recently Falco et al., (1995) demonstrated increased lysine in seeds of canola by 45% and in soybean by 33% by overexpressing
5 deregulated aspartate kinase and dihydrodipicolinate synthase. Since both the AK and DHPS enzymes were overexpressed relative to the wild-type enzymes, the biosynthetic pathway favored the formation of lysine over threonine (Figure 3). In all cases, the amount of available aspartate was not limiting in the formation of the end products.

10

In accordance with the present invention, threonine is an important metabolite in PHA biosynthesis as shown in Figure 3. Overexpression of either a wild-type or deregulated aspartate kinase will increase the available pools of free threonine in the plastids. As discussed in Examples
15 2-4, below, overexpression of a deregulated threonine deaminase in bacterial or plant cells will greatly enhance threonine turnover to α -ketobutyrate. A possible effect of this enhanced threonine turnover may be depletion of other aspartate family amino acids (see Figure 3). Increasing the level of aspartate kinase can ameliorate this possible
20 negative effect by increasing the pools of metabolic precursors involved in the biosynthesis of lysine, threonine, and methionine. This will result in enhancement of the free pools of lysine and methionine in the cell (i.e., in the plastids), thereby preventing starvation for lysine and/or methionine in the event that overexpression of threonine deaminase causes such
25 starvation. In addition to aspartate kinase (AK), homoserine dehydrogenase (HSD) and threonine synthase can be used to increase further the levels of free threonine (see Figure 3).

Deregulated aspartate kinases useful in the present invention can
30 possess a level of threonine and/or lysine insensitivity such that at 0.1 mM threonine and/or 0.1 mM lysine and the K_m concentration of aspartate, the

enzymes exhibit $\geq 10\%$ activity relative to assay conditions in which threonine and/or lysine is absent. Deregulated homoserine dehydrogenases useful in the present invention preferably possess a level of threonine insensitivity such that at 0.1 mM threonine and the K_m concentration of aspartate semialdehyde, the enzymes exhibit $\geq 10\%$ activity relative to assay conditions in which threonine is absent. The V_{max} values for the aspartate kinase and homoserine dehydrogenase enzymes can fall within the range of 0.1- 100 times that of their corresponding wild-type enzymes. The K_m values for the aspartate kinase and homoserine dehydrogenase enzymes can fall within the range of 0.01-10 times that of their corresponding wild-type enzymes.

Threonine synthase, the enzyme responsible for converting phosphohomoserine to threonine, has been shown to enhance the level of threonine about 10-fold over the endogenous level when overexpressed in *Methylobacillus glycogenes* (Motoyama et al., 1994). In addition, *E. coli* threonine synthase overexpressed in tobacco cell culture resulted in a 10-fold enhanced level of threonine from a six-fold increase in total threonine synthase activity (Muhitch, 1995). It is apparent from these results that overexpressed levels of threonine synthase in plants or other organisms will have the effect of increasing threonine levels therein. This can be employed in the present invention to insure an enhanced supply of threonine for α -ketobutyrate and propionyl-CoA production, the latter by the action of the next enzyme, threonine deaminase (described in Examples 2, 3, 4.). In conjunction with the PHA biosynthetic enzymes described herein, threonine synthase can enhance the formation of P(3HB-co-3HV) copolymer.

Example 2

Increased Production of α -Ketobutyrate From L-Threonine Via Modified *E. coli* Biosynthetic Threonine Deaminases

L-threonine can be degraded by several metabolic routes (Figure 3). Threonine aldolase (E.C. 4.1.2.5), a pyridoxal phosphate-containing enzyme, catalyzes threonine degradation to produce glycine and acetaldehyde, both of which can be metabolized to acetyl-CoA (Marcus and Dekker, 1993). Acetyl-CoA is the precursor of the C₄ monomer in PHB (Figure 1). L-threonine 3-dehydrogenase (E.C. 1.1.1.103) converts threonine into aminoacetone and CO₂ (Boylan and Dekker, 1981).

Alternatively, L-threonine can be converted to α -ketobutyrate by threonine deaminase (also designated threonine dehydratase, E.C. 4.2.1.16), as shown in Figure 3. α -ketobutyrate is a direct metabolic precursor of propionyl-CoA (Figure 3). There is also competition for α -ketobutyrate in isoleucine biosynthesis (Figure 3), and in transamination to produce 2-aminobutyrate (Shaner & Singh, 1993).

15 **Biodegradative Threonine Deaminase**

Threonine deaminase exists in two forms. One form is a biodegradative enzyme, encoded by, for example, the *tdcB* gene of *E. coli*. Biodegradative threonine deaminase permits threonine to be used as a carbon source. The structural gene for this enzyme has been isolated from *E. coli* K-12 (Goss and Datta, 1985), and the nucleotide sequence has been determined (Datta et. al., 1987). The biodegradative form of threonine deaminase is allosterically activated by high levels of AMP (Shizuta and Hayashi, 1976), and is inactivated by glucose, pyruvate, and other metabolites (Feldman and Datta, 1975). Biodegradative threonine deaminase is not feedback-inhibited by isoleucine. The K_m of *E. coli* biodegradative threonine deaminase for threonine has been determined to be 11 mM in the presence of activator (10 mM AMP) and 91 mM in its absence (Shizuta and Tokushige, 1971). The V_{max} for the enzyme decreases by a factor of six in the absence of activators (Shizuta and

Tokushige, 1971). Due to their requirement for activation by AMP to exhibit optimal activity *in vivo*, most degradative threonine deaminases may not be optimal for purposes of the present invention.

5 **Biosynthetic Threonine Deaminase**

Another form of threonine deaminase is a biosynthetic enzyme that catalyzes the committed step in isoleucine biosynthesis (Figure 3). In *E. coli*, this enzyme is encoded by the *ilvA* gene. It has recently been reported
10 (Colón et. al., 1995) that overexpression of the biosynthetic threonine deaminase from *C. glutamicum* in *C. lactofermentum* leads to an increase in the amount of isoleucine produced in the latter. In most cases reported, the goal of utilizing an overexpressed threonine deaminase was to increase the available pools of the essential amino acid isoleucine.

15

Biosynthetic threonine deaminase was the first enzyme reported to be feedback inhibited by the endproduct of its biosynthetic pathway (i.e., isoleucine), and served as a model for the development of the concept of allosteric regulation (Changeux, 1961; Monod et. al., 1963). Biosynthetic
20 threonine deaminase is strongly inhibited by isoleucine, and may not be optimal for producing enhanced levels of α -ketobutyrate in its natural form, even when overexpressed in recombinant systems. The results reported in Table 3, *infra*, support this conclusion. These results are discussed in greater detail in the section entitled "Biochemical Analysis of Wild-type and
25 Mutant *E. coli* Threonine Deaminases", below.

In addition to being inhibited by isoleucine, biosynthetic threonine deaminase exhibits kinetic positive cooperativity. This results in much slower substrate turnover at low concentrations of threonine as compared
30 to an enzyme that behaves in a normal Michaelis-Menten manner (also discussed in detail in the section entitled "Biochemical Analysis of Wild-type

and Mutant *E. coli* Threonine Deaminases", below). This too is an undesirable property for a threonine deaminase employed to produce high levels of α -ketobutyrate.

5 Isoleucine Deregulated Mutants of Biosynthetic Threonine Deaminase

Isoleucine-deregulated mutants of biosynthetic threonine deaminase have been described in the literature in both bacteria and
10 plants. One mutant, isolated from *S. typhimurium*, is referred to as *ilvA219* (LaRossa et. al., 1987). The *ilvA219* mutation results from a change in amino acid 447 from leucine (L) to phenylalanine (F). Another
isoleucine-deregulated mutant of biosynthetic threonine deaminase, identified in *E. coli*, is referred to as *ilvA466* (Taillon et al., 1988). The
15 *ilvA466* mutation results from a change in amino acid 481 from leucine (L) to phenylalanine (F). A third isoleucine-deregulated mutant of threonine deaminase has been described in *C. glutamicum*, and results from a change in amino acid 323 from valine (V) to alanine (A) (Möckel et al., 1994).

20 There have been two published reports of isoleucine-deregulated threonine deaminases in plants (Strauss et al., 1985; Mourad and King, 1995). In neither case was the mutation in the threonine deaminase identified.

25 Cloning and Expression of the Wild-type *E. coli* Biosynthetic Threonine Deaminase Gene

Conventional molecular biological techniques for routine cloning, transformation, expression, etc., are well known to those of ordinary skill in
30 the art and are described in detail in Sambrook et al., 1989 and Ausubel et al., 1989.

The nucleotide sequence of the wild-type *E. coli ilvA* threonine deaminase gene is available in the Genbank database (accession number K03503), and has also been published by Lawther et al. (1987). This
5 sequence is shown in SEQ ID NO:1.

In order to produce various isoleucine-deregulated mutant forms of *E. coli* threonine deaminase, the encoding genomic DNA was first isolated from *E. coli* DH5 α . Primers having the sequences shown in SEQ ID NO:2
10 and SEQ ID NO:3, respectively, were synthesized for use in the polymerase chain reaction (PCR) to amplify the gene encoding *E. coli* biosynthetic threonine deaminase:

SEQ ID NO:2

15

TTTTTGGATCCGATATCTTAACCCGCCAAAAAGAACCTGAACGCCG

SEQ ID NO:3

20 TTTTTGGATCCATGGCTGACTCGCAACCCCTGTCCGG

The 1,573 base pair PCR fragment thus obtained was subcloned as a BamHI restriction fragment into BamHI-digested plasmid pSP72 (Promega), creating pMON25659 (Figure 4). pMON25659 was digested
25 with NcoI, BstXI, and BamHI, and the two fragments containing the *ilvA* gene were purified and cloned as a triple ligation into NcoI and BamHI-digested pSE280 (Invitrogen), creating plasmid pMON25660 (Figure 5). Plasmid pMON25660 contains the IPTG-inducible ptrc promoter fused to the wild-type *E. coli* biosynthetic threonine deaminase gene (*ilvA*) for
30 expression in *E. coli*.

IlvA was expressed from plasmid pMON25660 by growing *E. coli* cells transformed with pMON25660 by CaCl₂/heatshock treatment in Luria Broth (LB) at 37°C to early log phase and inducing the cells with 0.5mM IPTG. The addition of IPTG to the cells results in high level
5 transcriptional activity from the ptrc promoter and expression of the IlvA.

IlvA protein extracts were prepared by pelleting induced cells, resuspending in 50mM KPi and 5% glycerol, sonicating, and removing cellular debris by centrifugation. Extracts of *E. coli* containing IlvA were
10 employed for biochemical studies as discussed below in the section entitled "Biochemical Analysis of Wild-type and Mutant *E. coli* Threonine Deaminases."

IlvA was further purified by the addition of protamine sulfate to the
15 protein extract (3% final concentration) and incubation on ice for one hour. The precipitate was removed by centrifugation and discarded. The supernatant was brought to 20% ammonium sulfate concentration, incubated at 4°C for one hour, and the precipitate was removed by centrifugation and discarded. Ammonium sulfate was added to the
20 supernatant to 50% final concentration, and the precipitate was pelleted by centrifugation. The pellet was resuspended in Buffer A: 10mM Bis-Tris-Propane, pH7.0, 1mM EDTA, 1mM DTT, and 1mM isoleucine. The protein solution was dialyzed (molecular weight cutoff of 12-14K) overnight in Buffer A. The dialyzed solution was loaded onto a Q-sepharose fast flow
25 column (2.5 cm x 40 cm, Pharmacia) pre-equilibrated with Buffer A. Elution of threonine deaminase was performed using 1250 ml Buffer A and 750 ml Buffer A plus 0.5M KCl (Buffer B). The elution gradient was increased from 0% to 75% B over 200 min. The flow rate was 10ml/min and fractions were collected every 2.5 min. Active fractions were pooled and the
30 enzyme precipitated by bringing the protein solution to 80% ammonium sulfate saturation. Precipitated protein was pelleted by centrifugation, the

pellet was dissolved in a small amount of Buffer A, and dialyzed overnight against Buffer A plus 0.01mM pyridoxal phosphate. The dialyzed protein solution was loaded onto a Mono-Q 10/10 column pre-equilibrated with Buffer A. Elution of threonine deaminase was performed using 400 ml Buffer A and 200 ml Buffer B in a gradient from 0% to 70% B over 140 min. The flow rate was 4 ml/min and 2 min fractions were collected. Active fractions with the highest specific activity were analyzed by SDS-PAGE/Coomassie Blue staining (data not shown). The purest fractions were pooled and used for the production of polyclonal antibodies in rabbits.

10 The remainder of the protein was stored at -80°C.

Mutagenesis of the *E. coli* Biosynthetic Threonine Deaminase Gene

Single Replacement of Leucine With Phenylalanine at Amino Acid Position 447

The cloned wild-type *E. coli* threonine deaminase gene (*ilvA*) (SEQ ID NO:1) was modified by substituting a leucine (L) codon for a phenylalanine (F) codon at amino acid position 447 as in the *S. typhimurium* *ilvA*219 enzyme (LaRossa et al., 1987) as follows.

The cloned *E. coli* *ilvA* threonine deaminase gene sequence in plasmid pMON25659 was excised as a BamHI fragment and cloned into BamHI- digested pGEM-3zf(-) (Promega Corp.), creating pMON25676 (Figure 6). Plasmid pMON25676 contains the M13 origin of replication for generation of single strand DNA (ssDNA). Single strand DNA from plasmid pMON25676 was prepared according to the pGEM® Single Strand systems procedure (Promega). The leucine codon corresponding to amino acid position 447 was modified by site-directed mutagenesis using BioRad's Muta-Gene® phagemid *in vitro* mutagenesis protocol employing the following synthetic mutagenic oligonucleotide (SEQ ID NO:4):

SEQ ID NO:4

5

CAGCTTCGAGTTCCCGGAATCACCGGGCGCGTTCCTGCGCTTCC

Specifically, the use of this synthetic mutagenic oligonucleotide modified SEQ ID NO:1 at nucleotide positions 1,317 (changing adenine to
10 guanine, therefore removing an EcoRI restriction enzyme site), 1,339
(changing cytosine to thymine), and 1,341 (changing guanine to cytosine).
The combined substitutions at nucleotides 1,339 and 1,341 changed the
codon for amino acid 447 from one encoding leucine to one encoding
phenylalanine. The mutagenesis successfully recreated the *S.*
15 *typhimurium ilvA219* mutation in the *E. coli* biosynthetic threonine
deaminase. The removal of the EcoRI restriction enzyme site allowed for
rapid screening of the resultant transformants following mutagenesis.
Successful mutagenesis was confirmed by testing the putative mutants for
insensitivity to feedback inhibition by isoleucine, as discussed below in the
20 section entitled "Biochemical Analysis of Wild-type and Mutant *E. coli*
Threonine Deaminases."

The nucleotide sequence of the mutagenized *E. coli* biosynthetic
threonine deaminase gene containing the *ilvA219* (L447F) mutation is
25 shown in SEQ ID NO:5. This mutant gene is contained in plasmid
pMON25679 (Figure 7).

**Single Replacement of Leucine With Phenylalanine at Amino
Acid Position 481**

30

The cloned wild-type *E. coli* threonine deaminase gene in pMON25676 was modified by site-directed mutagenesis by substituting a leucine (L) codon for a phenylalanine (F) codon at amino acid position 481 as in the *E. coli* IlvA466 enzyme (Taillon et al., 1988) using the mutagenic
 5 oligonucleotide of SEQ ID NO:6 as follows.

SEQ ID NO:6

TATCGCAGCCACGGCACCGACTACGGGCGCGTACTGGCGGCGTTTCG
 10 AATTTGGCGACCATGAACC

Use of this synthetic mutagenic oligonucleotide changed the codon for amino acid 481 from one encoding leucine to one encoding phenylalanine by changing nucleotide 1,441 of SEQ ID NO:1 from cytosine to thymine.
 15 This substitution successfully recreated the *E. coli* ilvA466 (L481F) mutation. This mutagenic oligonucleotide also changed nucleotide position 1,404 from thymine (T) to cytosine (C), thereby removing an NcoI restriction enzyme recognition site. The removal of the NcoI restriction enzyme site allowed for rapid screening of the resultant transformants
 20 following mutagenesis. Successful mutagenesis was confirmed by testing the putative mutants for insensitivity to feedback inhibition by isoleucine as discussed below in the section entitled "Biochemical Analysis of Wild-type and Mutant *E. coli* Threonine Deaminases."

25 The nucleotide sequence of the mutagenized *E. coli* biosynthetic threonine deaminase gene containing the *ilvA466* (L481F) mutation is shown in SEQ ID NO:7. This mutant gene is contained in plasmid pMON25680 (Figure 8).

30 Double Replacement of Leucine With Phenylalanine at Amino Acid Positions 447 and 481

Synthetic mutagenic oligonucleotides SEQ ID NO:4 and SEQ ID NO: 6 were used in concert to create a double *E. coli* mutant gene (L447F and L481F), referred to as *ilvA219/466*. The nucleotide sequence of this double mutant gene is shown in SEQ ID NO:8. The removal of the NcoI and EcoRI restriction enzyme sites allowed for rapid screening of the resultant transformants following mutagenesis. Successful mutagenesis was confirmed by testing the putative mutants for insensitivity to feedback inhibition by isoleucine as discussed below in the section entitled “Biochemical Analysis of Wild-type and Mutant *E. coli* Threonine Deaminases.” The *IlvA219/466* double mutant gene is contained in pMON25681 (Figure 9).

Cloning and Expression of the Mutant Threonine Deaminase Genes in *E. coli*

The plasmids pMON25679, pMON25680, and pMON25681 were separately digested with BstXI and BamHI, and the three mutant threonine deaminase genes obtained thereby were cloned into BstXI and BamHI-digested pMON25660 (Figure 5). The resultant *E. coli* expression plasmids contained the ptrc promoter fused to the mutant *ilvA* genes for overexpression in *E. coli*. The cloning resulted in the following plasmids: pMON25682 (Figure 10), containing the *ilvA219* mutation (L447F); pMON25683 (Figure 11), containing the *ilvA466* mutation (L481F); and pMON25684 (Figure 12), containing the *ilvA219/466* mutations (L447F/L481F). Expression of all the *IlvA* mutant enzymes was performed as described above in “Cloning and Expression of the Wild-type *E. coli* Biosynthetic Threonine Deaminase.” Extracts of *E. coli* containing the overexpressed mutant *IlvAs* were assayed for insensitivity to feedback inhibition by isoleucine as discussed below in the section entitled

"Biochemical Analysis of Wild-type and Mutant *E. coli* Threonine Deaminases."

Biochemical Analysis of Wild-type and Mutant Threonine Deaminases

5

Kinetic analysis of the wild-type and mutant threonine deaminases was performed by determining threonine turnover catalyzed by these threonine deaminases as assayed by the method of Burns (1971).

Specifically, α -keto-butyrate production was monitored by coupling to L-

10 lactic dehydrogenase (LDH)-catalyzed reduction of the α -ketoacid in the presence of NADH to form (L)-2-hydroxybutyrate and NAD⁺. The disappearance of NADH absorbance at 340 nm provides a measure of the threonine deaminase activity. Final reaction conditions were 100 mM KPi, pH 7.8, 0.20 mM NADH, 10 units/mL LDH (defined as LDH turnover of α -
15 ketobutyrate, not pyruvate), and variable threonine. *R. rubrum* was grown according to Brandl et al., 1989. The growth and induction of *E. coli* cells overexpressing the mutant and wild-type biosynthetic threonine deaminases is described above. Cell extracts of *R. rubrum* and *E. coli* were prepared by sonicating pelleted cells (\approx 0.5 g from 100 ml of culture) in 2 ml
20 of 100 mM KPi, pH 7.0, containing 5% (v/v) glycerol. After centrifugation to remove cellular debris, protein concentrations were determined by the method of Bradford (1976), and threonine deaminase activity was measured.

25 The kinetic parameters K_m and V_{max} , and the level of positive cooperativity, i.e., the deviation from normal saturation kinetics where at low substrate concentration the substrate turnover rate is less than a non-cooperative system, were determined. These values were found by fitting the rate data to the Hill equation (Equation 1), which incorporates the
30 coefficient n into the normal Michaelis-Menten equation ($n=1$ for a non-

cooperative system). An enzyme that displays no positive cooperativity exhibits an n value of ≤ 1 .

$$v = V_{\max} S^n / (K_m + S)^n \quad (1)$$

- 5 In addition to the kinetic parameters, the degree of inhibition of threonine deaminase by isoleucine was also measured at fixed levels of threonine.

The results are shown in Tables 1 and 2.

10

TABLE 1
General Rate Effects of Isoleucine on
Mutant and Wild-Type Threonine Deaminases

| Threonine deaminase | Threonine Deaminase Specific Activity, u/mg (%) ¹ | | |
|---|--|-------------------|--------------------|
| | 0 mM Isoleucine | 1.0 mM Isoleucine | 10.0 mM Isoleucine |
| <i>E. coli</i> wild-type | 16.7 | 0.3 (< 2) | 0.3 (< 2) |
| <i>E. coli</i> L447F | 16.0 | 16.3 (102) | 14.9 (93) |
| <i>E. coli</i> L481F | 20.3 | 11.3 (56) | 0.3 (< 2) |
| <i>E. coli</i> L447F/L481F | 22.0 | 22.8 (103) | 21.9 (99) |
| <i>R. rubrum</i> wild-type | 11.2 | no data | 10.0 (90) |
| <i>R. rubrum</i> wild-type (0.5 mM threonine) | 1.2 | no data | 1.1 (90) |

- 15 ¹ Rates were determined using 10 mM threonine as substrate, except where indicated, at the indicated concentrations of isoleucine. Values in parenthesis are % activity relative to the results with 0 mM isoleucine.

20 As shown in Table 1, the *E. coli* IlvA219 enzyme (*E. coli* L447F mutant) is insensitive to isoleucine inhibition up to a concentration of 10 mM isoleucine. The *E. coli* IlvA466 enzyme (*E. coli* L481F mutant) exhibits a more modest level of insensitivity to inhibition by isoleucine,

exhibiting 44% inhibition of enzyme activity at 1 mM isoleucine and 98% inhibition at 10 mM isoleucine. The novel double mutant enzyme (*E. coli* L447F/L481F mutant) is an improvement over the L447F mutant, exhibiting no inhibition at all in the presence of 10 mM isoleucine.

5

In addition to greatly enhanced insensitivity to isoleucine, the three mutant enzymes exhibit other enhanced kinetic properties. As shown in Table 2, the three mutant threonine deaminases exhibit lower K_m values and reduced or eliminated positive cooperativity compared to the wild-type enzyme. The latter is indicated by n values less than that of wild-type *E. coli* threonine deaminase (Table 2). The combination of a reduced n value and lower K_m can have a profound effect on the rate of substrate turnover, especially at low substrate concentrations. For example, using the reported K_m and n values for the threonine deaminases reported in Table 2, with equivalent V_{max} values and threonine = 0.5 mM, the L447F mutant is approximately a factor of 10 faster in substrate turnover than the wild-type enzyme.

20

TABLE 2
Kinetic Parameters for Mutant and Wild-type
Threonine Deaminases

| Enzyme ¹ | K_m Threonine (mM) | Hill Coefficient (n) |
|----------------------------|----------------------|--------------------------|
| <i>E. coli</i> wild-type | 5 ± 1 | 1.5 ± 0.2 |
| <i>E. coli</i> L447F | 2.01 ± 0.06 | 0.91 ± 0.03 |
| <i>E. coli</i> L481F | 3.08 ± 0.03 | 1.22 ± 0.01 |
| <i>E. coli</i> L447F/L481F | 2.15 ± 0.07 | 0.91 ± 0.04 |
| <i>R. rubrum</i> wild-type | 8.6 ± 0.3 | 0.79 ± 0.03 |

¹ The V_{max} for all threonine deaminases are within a factor of two of each other, being approximately 250-500 units/mg.

25

In vivo Analysis of Cloned *E. coli* *ilvA* Genes

5

In order to determine if overexpression of the wild-type *E. coli* biosynthetic threonine deaminase (IlvA) or an isoleucine-deregulated threonine deaminase mutant (for example, the IlvA466 (L481F) mutant *E. coli* threonine deaminase) could produce enhanced α -ketobutyrate levels or

- 10 P(3HB-co-3HV) copolymer from glucose and threonine in combination with the PHB biosynthetic operon of *A. eutrophus*, the following experiment was performed.

- E. coli* DH5 α cells were transformed with pJM9238 (Kidwell et al., 15 1995) containing the PHB operon of *A. eutrophus* by CaCl₂/heat shock treatment. The PHB operon was under the inducible control of the *ptac* promoter. *E. coli* containing pJM9236 were used as a background control with respect to threonine deaminase activity.

- 20 *E. coli* DH5 α cells were also cotransformed by CaCl₂/heat shock treatment with pJM9238 and pMON25660 (Figure 5) containing the wild-type *E. coli* threonine deaminase, under *ptrc* inducible control.

- In a third experiment, *E. coli* DH5 α cells were cotransformed with 25 pJM9238 and pMON25683 (Figure 11) containing the mutant *E. coli* *ilvA*466 (L481F) threonine deaminase, under *ptrc* inducible control.

- Transformed cells were grown on M9 minimal salts plus 0.2% gluconate and 25 mM L-threonine. At early log phase, the cells were 30 induced with IPTG to induce expression of all four enzymes (i.e., PhbA,

PhbB, PhbC, and IlvA or IlvA466). Threonine deaminase activity was measured in cell extracts as described above in the section entitled "Biochemical Analysis of Wild-type and Mutant *E. coli* Threonine Deaminases." Intracellular α -ketobutyrate was isolated from

5 approximately 50 mg (wet weight) of cells pelleted by centrifugation and then resuspended and sonicated in 1 ml of 100 mM KPi, pH 7.0, containing 5% glycerol (v/v). The level of α -ketobutyrate was determined by reverse-phase HPLC isolation of the dinitrophenylhydrazine derivative of the α -ketoacid according to Qureshi et al., (1982). Quantitation was based on

10 peak integration comparisons to a standard curve.

The PHA produced by the cells was extracted, hydrolyzed to the methylester, and analyzed by gas chromatography (GC) according to the following procedure. The polymer was first extracted by centrifuging the

15 cells at 7,000 rpm for 20 min., washing with 25 ml methanol, recentrifuging, washing with 15 ml hexane, and recentrifuging. Cell pellets were dried under N₂ for two hours, and the dry cell weights determined. 6.5 ml of chloroform were added to extract the PHA at 100°C for one hour. The solution was cooled and filtered through a PTFE syringe filter (13mm

20 diameter, 0.45 μ m pore). The PHA was precipitated by the addition of 50 ml of methanol, collected by centrifugation, and then washed with hexane. The polymer was dried at 70°C for two hours, and after drying, the polymer weight was determined.

25 The recovered PHAs were subjected to methanolysis by the method described by Brandl et al., (1988), with some modifications. Three to four milligrams of the polymer were dissolved in one ml chloroform containing a known amount (3.157 μ mol/ml in a typical example) of methyl benzoate internal standard (Aldrich Chemical Company, USA) in a small screw-cap

30 test tube. To this, 0.85 ml of methanol and 0.15 ml of concentrated sulfuric

acid were added, and the mixture was refluxed at 100°C for 140 min. At the end of the reaction, 0.5 ml of deionized water was added and the tube vortexed for one min. After phase separation, the organic phase (bottom layer), which contains the methyl esters of the constituent β -hydroxy-
5 carboxylic acids from the PHAs, was removed and dried over sodium sulfate, and then used for analysis.

The methyl esters were assayed by gas chromatography in order to determine the original copolymer composition, using a Varian 3400 gas
10 chromatograph equipped with a DB-5 capillary column (0.25 mm x 30 m, 0.25 μ film thickness; J&W Scientific, USA), a flame ionization detector, a Varian 8200 Autosampler, and a workstation with the Varian Star Chromatography software. 1 μ l of the sample was injected by split injection (split ratio of 50:1), using helium as the carrier gas (flow rate 2.5
15 ml/min). The temperatures of the injector and detector were kept at 250°C. A temperature program was used which separated the β -hydroxybutyric acid and β -hydroxyvaleric acid methyl esters efficiently (60°C for 3 min; temperature ramp of 12°C/min to 120°C; temperature ramp of 30°C/min to 300°C; 300°C for 10 min). Under these conditions,
20 the retention times of the methyl esters of β -hydroxybutyric acid and β -hydroxyvaleric acid were approximately 3.1 min and 4.8 min, respectively, while that of the methyl benzoate was 7.02 min. Quantitation was carried out based on standard curves generated using methyl-3-(D)-hydroxybutyrate and methyl-3-(D)-hydroxyvalerate (both obtained from
25 Fluka, USA) as external calibration standards. In order to take into account the partitioning (Jan et al., 1995) of the methyl esters between the organic and aqueous phases, as well as to correct for the differences in partition coefficients of the respective methyl esters, the standards were treated in a manner identical to that described above for the PHAs (except
30 for the omission of the reflux step) prior to generating the standard curves.

The mol % C5 (3HV component) was determined in the isolated polymer. The results are shown in Table 3.

TABLE 3

5 **Effect of Threonine Deaminase on α -Ketobutyrate
Production and %C5 Content in PHBV³**

| Tda ¹ <i>E. coli</i> construct | Threonine Deaminase Specific Activity, u/mg (%) ² | | [α -ketobutyrate] μ M | % PHBV Dry Weight (%C5) |
|--|---|----------------------|--------------------------------------|----------------------------------|
| | 0 mM Isoleucine | 0.1 mM Isoleucine | | |
| pJM9238 ⁴ | < 0.04 | < 0.04 | 13.4 | 35 (1.3) |
| pJM9238/pMON25660 ⁵ | 4.20 | 0.16 (3.8) | 41.8 | 22 (1.1) |
| pJM9238/pMON25683 ⁶ | 0.59 | 0.56 (95) | 579 | 27 (0.9) |

¹ Tda refers to threonine deaminase.

2 Rates were determined using 10 mM threonine as substrate, at the
10 indicated concentrations of isoleucine. Values in parenthesis are % activity
relative to the 0 mM isoleucine results.

³ PHBV refers to P(3HB-co-3HV) copolymer.

⁴ pJM9238 contains the *A. eutrophus* PHB operon.

⁵ pMON25660 contains the wild-type *E. coli* threonine deaminase.

15 ⁶ pMON25683 contains the L481F mutant *E. coli* threonine deaminase.

Table 3 shows that there is about a three-fold increase in α -ketobutyrate concentration for the overexpressed wild-type threonine deaminase compared to cells not overexpressing a threonine deaminase
20 (41.8 μ M vs. 13.4 μ M α -keto-butyrate). This is evidently a reflection of the strong feedback regulation of the wild-type enzyme by isoleucine *in vivo*. In contrast, the transformed cells containing the L481F mutant threonine

deaminase accumulated α -keto-butyrate to a level greater than 40 times that in cells containing no overexpressed threonine deaminase (579 μ M vs. 13.4 μ M). This occurred despite the fact that the threonine deaminase activity in the transformants containing overexpressed wild-type enzyme was much greater than that in transformants containing the L481F mutant (4.20 u/mg vs. 0.59 u/mg in the absence of isoleucine). It is also noteworthy that the α -ketobutyrate concentrations reported in Table 3 are for a 1 ml extract solution from approximately 50mg of cells, and therefore represent a significant dilution of the intracellular metabolite, probably by a minimum factor of 10.

If simply increasing the levels of α -ketobutyrate was sufficient to produce a significant 3HV component in the P(3HB-co-3HV) copolymer, then there should have been significantly more 3HV in the copolymer when employing pJM9238/pMON25683 than when employing pJM9238/pMON25660. As shown in Table 3, this was not the case. This result suggests either that the α -ketobutyrate was not being converted to propionyl-CoA, or that the *A. eutrophus* PHB biosynthetic enzymes encoded by pJM9238 could not efficiently catalyze formation of the C5 substrate ((D)-3-hydroxy-valeryl-CoA, Figure 2) for incorporation into the polymer. The results presented in Example 8 strongly suggest that this is at least partially due to a block in the ability of the *A. eutrophus* PhbA to catalyze the condensation of acetyl-CoA and propionyl-CoA.

In summary, it is evident from the data in Tables 1-3 that mutant, deregulated threonine deaminases similar to those described herein possess a desirable deregulated phenotypic property useful in enhancing the level of α -ketobutyrate in organisms such as bacteria and plants targeted for the production of P(3HB-co-3HV). Isoleucine-deregulated threonine deaminases useful in the present invention preferably possess a level of

isoleucine insensitivity such that at 100 μ M isoleucine and 10 mM threonine, the enzymes exhibit $\geq 10\%$ activity relative to assay conditions in which isoleucine is absent. In addition to exhibiting reduced isoleucine sensitivity, threonine deaminases useful in the present invention can also exhibit reduced or no positive cooperativity compared to the corresponding wild-type enzymes (Hill coefficient < 1.5), and have kinetic parameters of K_m and V_{max} in the range of the wild-type enzyme (e. g., $V_{max} = 10$ -1000U/mg, $K_m = 0.1$ to 20mM).

10 The present invention encompasses not only the DNA sequences shown in SEQ ID NOS:5,7, and 8 and the proteins encoded thereby, but also biologically functional equivalent nucleotide and amino acid sequences. The phrase "biologically functional equivalent nucleotide sequences" denotes DNAs and RNAs, including chromosomal DNA, plasmid DNA, 15 cDNA, synthetic DNA, and mRNA nucleotide sequences, that encode peptides, polypeptides, and proteins exhibiting the same or similar threonine deaminase enzymatic activity as that of the mutant *E. coli* threonine deaminases encoded by SEQ ID NOS:5, 7, and 8, respectively, when assayed enzymatically by the methods described herein, or by 20 complementation. Such biologically functional equivalent nucleotide sequences can encode peptides, polypeptides, and proteins that contain a region or moiety exhibiting sequence similarity to the corresponding region or moiety of the *E. coli* IlvA muteins.

25 One can isolate threonine deaminases useful in the present invention from various organisms based on homology or sequence identity. Although particular embodiments of nucleotide sequences encoding deregulated threonine deaminases are shown in SEQ ID NOS:5, 7, and 8, it should be understood that other biologically functional equivalent forms of 30 such deregulated threonine deaminase-encoding nucleic acids can be readily isolated using conventional DNA-DNA or DNA-RNA hybridization

techniques. Thus, the present invention also includes nucleotide sequences that hybridize to any of SEQ ID NOS:5, 7, and 8 and their complementary sequences, and that code on expression for peptides, polypeptides, and proteins exhibiting the same or similar enzymatic activity as that of these
5 deregulated threonine deaminases. Such nucleotide sequences preferably hybridize to SEQ ID NO:5, 7, or 8 or their complementary sequences under moderate to high stringency (see Sambrook et al., 1989). Exemplary conditions include initial hybridization in 6X SSC, 5X Denhardt's solution, 100 µg/ml fish sperm DNA, 0.1% SDS, at 55°C for sufficient time to permit
10 hybridization (e.g., several hours to overnight), followed by washing two times for 15 min each in 2X SSC, 0.1% SDS, at room temperature, and two times for 15 min each in 0.5-1X SSC, 0.1% SDS, at 55°C, followed by autoradiography. Typically, the nucleic acid molecule is capable of hybridizing when the hybridization mixture is washed at least one time in
15 0.1X SSC at 55°C, preferably at 60°C, and more preferably at 65°C.

The present invention also encompasses nucleotide sequences that hybridize under salt and temperature conditions equivalent to those described above to genomic DNA, plasmid DNA, cDNA, or synthetic DNA
20 molecules that encode the same amino acid sequences as these nucleotide sequences, and genetically degenerate forms thereof due to the degeneracy of the genetic code, and that code on expression for a peptide, polypeptide, or protein that has the same or similar threonine deaminase enzymatic activity as that of the deregulated threonine deaminases disclosed herein.

25

Biologically functional equivalent nucleotide sequences of the present invention also include nucleotide sequences that encode conservative amino acid changes within the amino acid sequences of the present deregulated threonine deaminases, producing silent changes
30 therein. Such nucleotide sequences thus contain corresponding base

substitutions based upon the genetic code compared to the nucleotide sequences encoding the present deregulated threonine deaminases.

Substitutes for an amino acid within the fundamental mutant deregulated *E. coli* threonine deaminase amino acid sequences discussed herein can be

- 5 selected from other members of the class to which the naturally occurring amino acid belongs. Amino acids can be divided into the following four groups: (1) acidic amino acids; (2) basic amino acids; (3) neutral polar amino acids; and (4) neutral non-polar amino acids. Representative amino acids within these various groups include, but are not limited to: (1) acidic
- 10 (negatively charged) amino acids such as aspartic acid and glutamic acid; (2) basic (positively charged) amino acids such as arginine, histidine, and lysine; (3) neutral polar amino acids such as glycine, serine, threonine, cyteine, cystine, tyrosine, asparagine, and glutamine; and (4) neutral nonpolar (hydrophobic) amino acids such as alanine, leucine, isoleucine,
- 15 valine, proline, phenylalanine, tryptophan, and methionine.

- Conservative amino acid changes within the present mutant deregulated *E. coli* threonine deaminase sequences can be made by substituting one amino acid within one of these groups with another amino
- 20 acid within the same group. While biologically functional equivalents of these mutant deregulated threonine deaminases can have any number of conservative amino acid changes that do not significantly affect the threonine deaminase enzymatic activity of this enzyme, 10 or fewer conservative amino acid changes may be preferred. More preferably, seven
- 25 or fewer conservative amino acid changes may be preferred; most preferably, five or fewer conservative amino acid changes may be preferred. The encoding nucleotide sequences (gene, plasmid DNA, cDNA, synthetic DNA, or mRNA) will thus have corresponding base substitutions, permitting them to code on expression for the biologically functional
- 30 equivalent forms of the mutant deregulated *E. coli* threonine deaminases.

In addition to nucleotide sequences encoding conservative amino acid changes within the amino acid sequences of the present mutant threonine deaminases, biologically functional equivalent nucleotide sequences of the present invention also include genomic DNA, plasmid DNA, cDNA, synthetic DNA, and mRNA nucleotide sequences encoding non-conservative amino acid substitutions, additions, or deletions. These include nucleic acids that contain the same inherent genetic information as that contained in the DNA of SEQ ID NOS:5, 7, and 8, and which encode peptides, polypeptides, or proteins exhibiting the same or similar enzymatic activity as that of the present mutant deregulated threonine deaminases. Such nucleotide sequences can encode fragments or variants of these threonine deaminases. The enzymatic activity of such fragments and variants can be determined by complementation or enzymatic assays as described above. These biologically functional equivalent nucleotide sequences can possess from 40% sequence identity, or from 60% sequence identity, or from 80% sequence identity, to 100% sequence identity to the DNAs encoding the present deregulated threonine deaminases, or corresponding regions or moieties thereof. However, regardless of the percent sequence identity of these biologically functional equivalent nucleotide sequences, the encoded proteins would possess the same or similar enzymatic activity as the deregulated threonine deaminases disclosed herein. Thus, biologically functional equivalent nucleotide sequences encompassed by the present invention include sequences having less than 40% sequence identity to any of SEQ ID NOS:5, 7, and 8, so long as they encode peptides, polypeptides, or proteins having the same or similar enzymatic activity as the deregulated threonine deaminases disclosed herein.

Mutations made in *E. coli* threonine deaminase cDNA, chromosomal DNA, plasmid DNA, synthetic DNA, mRNA, or other nucleic acid preferably preserve the reading frame of the coding sequence.

Furthermore, these mutations preferably do not create complementary regions that could hybridize to produce secondary mRNA structures, such as loops or hairpins, that would adversely affect mRNA translation.

Useful biologically functional equivalent forms of the DNAs of SEQ

- 5 ID NOS:5, 7, and 8 include DNAs comprising nucleotide sequences that exhibit a level of sequence identity to corresponding regions or moieties of these DNA sequences from 40% sequence identity, or from 60% sequence identity, or from 80% sequence identity, to 100% sequence identity to the DNAs encoding the present deregulated threonine deaminases, or
- 10 corresponding regions or moieties thereof. However, regardless of the percent sequence identity of these nucleotide sequences, the encoded proteins would possess the same or similar enzymatic activity as the deregulated threonine deaminases disclosed herein. Thus, biologically functional equivalent nucleotide sequences encompassed by the present
- 15 invention include sequences having less than 40% sequence identity to any of SEQ ID NOS:5, 7, and 8, so long as they encode peptides, polypeptides, or proteins having the same or similar enzymatic activity as the deregulated threonine deaminases disclosed herein. Sequence identity can be determined using the "BestFit" or "Gap" programs of the Sequence
- 20 Analysis Software Package, Genetics Computer Group, Inc., University of Wisconsin Biotechnology Center, Madison, WI 53711.

- Due to the degeneracy of the genetic code, i.e., the existence of more than one codon for most of the amino acids naturally occurring in proteins,
- 25 genetically degenerate DNA (and RNA) sequences that contain the same essential genetic information as the DNAs of SEQ ID NOS:5, 7, and 8 of the present invention, and which encode the same amino acid sequences as these nucleotide sequences, are encompassed by the present invention. Genetically degenerate forms of any of the other nucleic acid sequences
- 30 discussed herein are encompassed by the present invention as well.

The nucleotide sequences described above are considered to possess a biological function substantially equivalent to that of the mutant *E. coli* threonine deaminase-encoding DNAs of the present invention if they encode peptides, polypeptides, or proteins having isoleucine-deregulated threonine deaminase enzymatic activity differing from that of any of the mutant *E. coli* threonine deaminases disclosed herein by about $\pm 30\%$ or less, preferably by about $\pm 20\%$ or less, and more preferably by about $\pm 10\%$ or less when assayed *in vivo* by complementation or by the enzymatic assays discussed above.

10

Example 3

Increased Production of α -Ketobutyrate Via Use of the Threonine Deaminase of *Rhodospirillum rubrum*

The phototrophic, non-sulfur purple bacterium *Rhodospirillum rubrum* (ATCC 25903) is the source of a natural deregulated threonine deaminase (Feldberg and Datta, 1971). As shown in Tables 1 and 2, above, the threonine deaminase from *R. rubrum* displays normal Michaelis-Menten behavior, and is not significantly feedback inhibited by isoleucine. This enzyme is also unaffected by activators of the degradative enzyme (Feldberg and Datta, 1971). The *R. rubrum* threonine deaminase can therefore also be expressed in bacteria and plants in order to enhance the production of α -ketobutyrate from L-threonine.

SDS-PAGE analysis indicates that the monomeric molecular weight of the *R. rubrum* threonine deaminase is approximately 42 kD (data not shown). This is intermediate between the molecular weight of the *E. coli* biosynthetic (56 kD) and biodegradative enzymes (35 kD). The *R. rubrum* enzyme may therefore represent a third class of threonine deaminase useful in the present invention. Möckel et al. (1994) similarly observed that

the threonine deaminase of *C. glutamicum* was of intermediate molecular weight, having a C-terminal deletion of 95 amino acids.

From what is known of other threonine deaminases, it is probable
5 that the C-terminal domain of the *R. rubrum* enzyme is truncated,
resulting in the deregulated phenotype of the enzyme. Taillon et al. (1988)
published an amino acid comparison of three biosynthetic threonine
deaminases and one biodegradative threonine deaminase. This comparison
revealed N-terminal conservation between the two forms of threonine
10 deaminase. The N-terminus appears to be involved in catalysis. The
biodegradative form of the enzyme is truncated at the C-terminus, and it
has been suggested that this deletion results in the isoleucine-deregulated
phenotype (Taillon et al., 1988). Based on SDS-PAGE analysis of the *R.*
rubrum threonine deaminase which indicates that the *R. rubrum* enzyme
15 possesses a molecular weight intermediate between that of the biosynthetic
and degradative enzymes, the *R. rubrum* threonine deaminase may have a
truncated C-terminus, effectively deleting the regulatory domain.

Example 4

Increased Production of α -Ketobutyrate Using Other Modified or Naturally-Occurring Forms of Threonine Deaminase

Deregulated threonine deaminases useful in the present invention
can be either naturally-occurring, or produced by recombinant DNA
25 techniques such as those employed in Example 2. If naturally-occurring,
such threonine deaminases can be identified by screening organisms for
threonine deaminase activity, followed by measuring the sensitivity of that
activity to isoleucine inhibition using the standard assay methods described
above. Genes for these enzymes can then be isolated by complementation
30 into threonine deaminase-negative bacteria (Fisher and Eisenstein, 1993).
The levels of isoleucine analogs such as L-O-methylthreonine (OMT) in the

growth medium can be varied during the complementation procedure to select organisms that express isoleucine-deregulated biosynthetic threonine deaminases. Alternatively, genes for threonine deaminases can be obtained by hybridization techniques using an appropriate probe,

- 5 preferably based upon the nucleotide sequence comprising the N-terminal region of the biosynthetic threonine deaminase. Following expression of these isolated genes in an appropriate host, they can be tested for isoleucine insensitivity as described previously.

- 10 Alternatively, deregulated threonine deaminases can be created by physical or chemical mutagenesis of wild-type threonine deaminase genes utilizing a variety of techniques known in the art to introduce either specific or random mutations into such genes (Eisenstadt et al., 1994; as described in Example 2; Feldmann et al., 1994). Muteins created in this way can be
15 analyzed by methods such as those described in Example 2 to determine catalytic efficiencies and the extent of inhibition by isoleucine.

- One strategy for creating deregulated threonine deaminases useful in the present invention is based upon knowledge of the functional domains
20 of this enzyme. The functional domains of both the degradative and biosynthetic types of threonine deaminase have been analyzed (Taillon et al., 1988). All known mutations affecting feedback inhibition of the biosynthetic threonine deaminase are C-terminal region mutations. If the biosynthetic threonine deaminase is compared to the biodegradative
25 threonine deaminase at the amino acid level, it can be seen that the regions of homology between the two forms exist only in the N-terminal regions of the polypeptides. Since the biodegradative form of the enzyme is not regulated by isoleucine, the N-terminal region is likely responsible for the catalytic activity, while the C-terminal region is likely involved in the
30 feedback regulation (Fisher and Eisenstein, 1993). This knowledge of the regulatory domains of the enzyme suggests that modification of the C-

terminal region will produce mutants with desirable properties for the purposes disclosed herein. For example, portions of the C-terminal region of the biosynthetic threonine deaminase can be sequentially deleted, and the remainder of the enzyme polypeptide assayed for activity to isolate

- 5 deregulated deletion mutants. Deletions of the C-terminal region of the biosynthetic deaminase can be produced by a variety of techniques known in the art. For example, Exonuclease III can be used to create a ladder of 3' deletions. Following expression of the truncated genes, the enzymatic activity of the modified deaminases can be assayed as described above.

10

Alternatively, various domains of different threonine deaminases can be replaced with one another to determine their effect on enzyme regulation. For example, the C-terminal region of a biosynthetic threonine deaminase can be replaced with the C-terminal region of a biodegradative

15 threonine deaminase such as that from *E. coli*, and the resulting polypeptide assayed for feedback inhibition by isoleucine. Domain swapping has been discussed in Cohen and Curran 1990; Czerny et. al., 1993; and Zinszner et. al., 1994.

20

Mutants of biodegradative threonine deaminase that are not inhibited by pyruvate, or which do not require activation by AMP, can be generated by random mutagenesis of a cloned biodegradative threonine deaminase, or by swapping domains of the biodegradative threonine deaminase with those of the biosynthetic enzyme. Such mutant

25

biodegradative threonine deaminases can be use to generate high levels of α -ketobutyrate in plants or bacteria. No biodegradative threonine deaminase has yet been reported that does not require a metabolite for activation.

30

Example 5**5 Expression and Activity of Wild-type and Mutant Threonine Deaminases in Plant Cells****Expression and Activity in Electroporated Tobacco Protoplasts**

10 In order to test the expression and enzymatic activity of the cloned *E. coli* biosynthetic threonine deaminases in plant cells, several plant expression plasmids were constructed for transient expression in tobacco leaf protoplasts. The plant expression plasmids contain all the necessary elements for plant cell expression, including a promoter (i.e., e35S (Odell et

15 al., 1985), the 5' untranslated leader sequence within the e35S promoter, a coding sequence, and a transcription termination and polyadenylation signal (i.e., E9 (Coruzzi et al., 1984)). The wild-type *E. coli* biosynthetic threonine deaminase (IlvA) encoded by the insert in pMON25663 (Figure 13) and the three mutant, deregulated *E. coli* biosynthetic threonine

20 deaminases encoded by the inserts in pMON25686 (Figure 14), pMON25687 (Figure 15), and pMON25688 (Figure 16) containing the IlvA 219 (L447F), IlvA466 (L481F), and IlvA219/466 (L447F/L481F) mutations, respectively, were cloned downstream of the e35S constitutive plant promoter (Odell et al., 1985) and targeted to chloroplasts using a

25 translational fusion to the *Arabidopsis* rubisco small subunit chloroplast transit peptide (ArabSSU1A; Stark et al., 1992). The four plasmids containing the various threonine deaminase-encoding DNAs were independently electroporated into tobacco leaf protoplasts by the method of Hinchey et al. (1994). Tobacco protoplasts electroporated with the various

30 DNA's were sonicated to lyse the protoplasts. Figure 17 shows the expression of the various IlvA enzymes determined by Western blot

analysis using rabbit polyclonal antibodies generated to the wild-type IlvA. Enzyme activity was monitored in extracts of lysed protoplasts in the presence and absence of isoleucine as described above, and is reported in Table 4.

5

TABLE 4
Activity of Threonine Deaminases
Introduced Into Tobacco Protoplasts

| Electroporated Plasmid | Threonine Deaminase Specific Activity, u/mg (%) ¹ | |
|--------------------------------------|--|----------------|
| | 0 mM Isoleucine | 1mM Isoleucine |
| no plasmid DNA | 0.000 | 0.000 |
| pMON25663 (wt IlvA) | 0.089 | 0.006 (6.7) |
| pMON25686 (IlvA 219 L447F) | 0.101 | 0.091 (90.1) |
| pMON25687 (IlvA 466 L481F) | 0.162 | 0.048 (29.6) |
| pMON25688 (IlvA 219/466 L447F/L481F) | 0.181 | 0.181 (100) |

10 ¹ Rates were determined using 10 mM threonine as substrate, at the indicated concentrations of isoleucine. Values in parenthesis are % activity relative to the results with 0 mM isoleucine.

15 As shown in Figure 17, the "no DNA" control tobacco protoplasts did not exhibit any immunoreactive bands corresponding to threonine deaminase by Western blotting. In contrast, extracts of lysed tobacco protoplasts into which plasmids pMON25663, pMON25686, pMON25687, and pMON25688 were electroporated contained immunoreactive bands
20 corresponding to the predicted molecular weight of the threonine deaminase encoded by the inserts of the respective plasmids. The enzymatic activity

data reported in Table 4 demonstrate that both wild-type and deregulated threonine deaminases exhibit their predicted isoleucine deregulated enzymatic activities when expressed in plant cells based upon the results reported in the section entitled "Biochemical Analysis of Wild-type and
5 Mutant *E. coli* Threonine Deaminases" of Example 2.

Expression and Activity in Transformed Soybean Callus

As shown above, the *E. coli* IlvA proteins can be functionally
10 expressed in a plant transient expression system. In order to test the expression of a biosynthetic threonine deaminase in a stably transformed plant tissue, non-differentiating callus derived from soybean hypocotyl tissue was transformed with a plasmid containing the *E. coli* wild-type *ilvA* threonine deaminase gene using *Agrobacterium tumefaciens* ABI (Koncz
15 and Schell, 1986).

Transformation Vector

Soybean callus tissue was transformed with binary Ti plasmid
20 pMON25668 (Figure 18). Plasmid pMON25668 contains the e35S-expressed, plastid-targeted (ARABSSU1A) *E. coli* wild-type *ilvA* threonine deaminase gene, the e35S-expressed, plastid-targeted (ARABSSU1A) *A. eutrophus phbA* (β -ketothiolase) gene, and the e35s-expressed neomycin phosphotransferase type II (NptII) gene for kanamycin resistance.
25 pMON26668 contains two border sequences for T-DNA transfer into the plant chromosome. The right border sequence is the only border required for T-DNA transfer into the plant; however, including the left border sequence terminates T-DNA integration at that point, thereby limiting the unnecessary incorporation of "extra" DNA sequences into the plant
30 chromosome. pMON25668 also contains the minimum sequence of ori-322 for replication and maintenance in *E. coli* as well as ori-V of the broad host

origin RK2 for replication in *Agrobacterium*. TrfA is supplied in trans for proper replication in *Agrobacterium*. Further details of these vectors can be found in Glick et al., (1993) and the references cited therein.

5 **Seedling Growth**

The bottom of a 100 X 25 mm Petri dish was covered with Asgrow A3237 soybean seeds and the lid was replaced. Four Petri dishes containing seeds were placed inside a vacuum bubble containing a glass
10 beaker holding 200 mls of Chlorox (sodium hypochlorite). The bubble was placed in a chemical fume hood. Two mls of HCl were added to the Chlorox, and the bubble lid was placed on quickly. A vacuum was pulled just long enough for the lid to stay on tightly, and the seeds were exposed to the resulting fumes overnight. The following day, the Petri dishes were removed
15 from the bubble, and dry Captan (1/8 tsp) was added to each dish. The seeds were covered with sterile water and incubated in the Captan slurry for 5-7 minutes, after which the Captan slurry was removed.

After pipetting off the liquid from all of the plates, the seeds (20-25
20 per plate) were placed on 0.8% water agar plates. Only good quality seeds were chosen, i.e., no seeds with cracked seedcoats, discoloration, or other damage were used. Five plate high stacks of Petri plates were then wrapped with a rubberband, covered with a clear plastic bag, and incubated in a warm room set at 25°C under continuous cool white light (60 $\mu\text{En m}^{-2}\text{s}^{-1}$;
25 this can range from about 20-80 $\mu\text{En m}^{-2}\text{s}^{-1}$) for 6 days to obtain seedlings.

Preparation of *Agrobacterium*

30 About one week prior to inoculation, *Agrobacterium tumefaciens* ABI was streaked from a glycerol stock onto LB agar-solidified plates (1.5%

agar) containing 100 mg/l spectinomycin, 50 mg/l kanamycin, and 25 mg/l chloramphenicol, and grown at room temperature.

The day before explant inoculation, *Agrobacterium* cultures were
5 started in clear plastic tubes containing 2 mls of YEP (Sambrook et al.,
1989) containing the same levels of spectinomycin, kanamycin, and
chloramphenicol as above. About 1/4 loopful of bacteria was placed in each
YEP tube. The tubes were placed on a rotator in a warm room set at 25°C.
Eight hours later, each 2 ml culture was added to 25 mls liquid YEP
10 containing the same amounts of antibiotics as above, as well as 200µM
acetosyringone and 1mM galacturonic acid, in a sterile 250 ml flask. The
cultures were grown overnight in the dark at 28°C with shaking (170 rpm).

On the day of explant inoculation, 12 ml aliquots of *Agrobacterium*
15 were placed in sterile 50 ml centrifuge tubes. The tubes were centrifuged
for 12 minutes at 2,000 rpm in order to pellet the cells. The supernatants
were poured off and the pellets were resuspended in 20 mls of co-culture
medium containing 1/10 MS salts and 1/10 B5 vitamins (Sigma, M 0404)
and 15 g/l glucose, 20mM MES, pH 5.4, and combined. The cell density
20 was adjusted to a final OD₆₆₀ in the range from 0.3-0.35 from an initial
OD₆₆₀ of 0.5-0.6. Thirty mls of this *Agrobacterium* suspension were used to
inoculate each batch of 50 hypocotyl explants.

Explant Inoculation and Co-culturing

25

One day prior to explant inoculation, the plates containing the 6 day
old seedlings were taken from the warm room and placed in a refrigerator at
0-10°C (average temperature of 4°C) for approximately 24 hrs. On the day
of explant inoculation, the stacks of seedlings were taken from the cold one
30 at a time just prior to explanting to maintain coldness. The hypocotyls
were cut into 5mm sections, and batches of 50 sections were inoculated by

incubating them with 30 mls of *Agrobacterium* suspension in a Petri plate (100X25 mm) for 30 min. After 30 min, the *Agrobacterium* suspension was pipetted off. The explants were blotted on sterile qualitative Whatman filter paper, and then placed 10 per plate in 100X15 mm Petri plates

- 5 containing one 8.5 cm sheet of sterile Whatman filter paper and 4 mls/plate of liquid co-culture medium containing 1/10 MS salts and 1/10 B5 vitamins, 15 g/l glucose, 3.9 g/l MES, 4.68 mg/l naphthaleneacetic acid (this can range from about 1-8 mg/l), 2.5 mg/l kinetin (this can range from about 0.5-4 mg/l), 200 μ M acetosyringone (this can range from about 50-300 μ M), and 1
10 mM galacturonic acid (this can range from about 0.1 - 2 mM), at pH 5.4. The plates were wrapped with parafilm and incubated for two days in a warm room set at 25°C under continuous cool white light (40 μ En m⁻²s⁻¹; this can range from about 20-60 μ En m⁻²s⁻¹).

15 Selection

- After the two day co-culture period, the explants were placed 10 per plate onto solid selection medium containing 1X MS salts and 1X B5 vitamins (Sigma, M 0404), 3% sucrose (this can range from about 1-6%),
20 4.68 mg/l naphthalene- acetic acid (this can range from about 1-8 mg/l), 2.15 mg/l kinetin (this can range from about 0.5-4 mg/l), 500 mg/l ticarcillin (this can range from about 25-250 mg/l) , 100 mg/l kanamycin (this can range from about 25-250 mg/l), and 0.7% purified agar (Sigma, B11853), at pH 5.7. The plates were wrapped with white 3M filter tape and cultured in
25 a warm room set at 25°C under continuous cool white light (40 μ En m⁻²s⁻¹; this can range from about 20-60 μ En m⁻²s⁻¹). Explants were transferred to fresh medium every two weeks. At the six week timepoint, the calli were excised from the hypocotyls and again transferred to the same selection medium for an additional two weeks. All calli were maintained on this
30 selection medium until they were ready for assaying at 10-16 weeks.

Kanamycin-resistant calli were analyzed by Western blotting and enzymatic assays as described above to determine if *E. coli* IlvA was expressed and exhibited enzymatic activity in stably transformed soybean cells. The results are shown in Figure 19.

5

The data in Figure 19 demonstrate that in four transformed calli (i.e., 518-4, 518-15, 518-16, and 518-17), the *E. coli* IlvA biosynthetic threonine deaminase was detectable by Western blot analysis. Enzymatic activity of the IlvA was also confirmed in the same transformed calli, correlating with the positive Western events. These results demonstrate that the *E. coli* IlvA can be expressed and maintain enzymatic activity in stably transformed plant tissue.

Colau et al., (1987) expressed the *S. cerevisiae* wild-type Ilv1 gene in tobacco (*N. plumbaginifolia*) mutants deficient in threonine deaminase. Expression of this threonine deaminase in these mutants was sufficient to complement the isoleucine auxotrophy.

Example 6

Increased Propionyl-CoA Production From α -Ketobutyrate Via Modification of the Pyruvate Dehydrogenase Complex or Expression of a Branched-Chain α -Ketoacid Dehydrogenase Complex

As demonstrated for the bacterial (Danchin et al., 1984; Bisswanger, 1981) and pea chloroplast complexes (Camp et al., 1988; Camp and Randall, 1985), pyruvate dehydrogenase complex (PDC) catalyzes the oxidative decarboxylation of α -ketobutyrate to produce propionyl-CoA (Figure 3). The rates of turnover of α -ketobutyrate with both the bacterial and chloroplast PDCs are approximately 10% of that

relative to pyruvate at 1.5 mM concentration of α -ketoacid. The effect on kinetic constants with the bacterial enzyme is a 10-fold increase in K_m , and a five-fold decrease in V_{max} (Bisswanger, 1981). Therefore, if steady-state concentrations of α -ketobutyrate rise to high μ M or low mM levels,

- 5 turnover of α -ketobutyrate to propionyl-CoA catalyzed by the pyruvate dehydrogenase complex should occur in plants and bacteria. Van Dyk and LaRossa (1987) demonstrated in wild-type *Salmonella typhimurium*, under conditions where acetohydroxyacid synthase (AHAS, Figure 3) activity is reduced by inhibition with sulfometuron methyl, that the concentration of
- 10 α -ketobutyrate increases significantly, and propionyl-CoA accumulates (acetohydroxyacid synthase condenses α -ketobutyrate and pyruvate to produce 2-aceto-2-hydroxy-isovalerate as the second step in the biosynthesis of isoleucine, see Figure 3). With respect to the P(3HB-co-3HV) copolymer biosynthetic pathway depicted in Figure 3 and as
- 15 discussed in Examples 2-4 above, high levels of α -ketobutyrate produced by the overexpression of a deregulated threonine deaminase should be sufficient for propionyl-CoA formation, even when there is competing demand for the α -ketoacid in isoleucine biosynthesis. Results presented in
- 20 Table 3 demonstrating elevated levels of α -ketobutyrate *in vivo* from the presence of a deregulated threonine deaminase support this hypothesis.

The pyruvate dehydrogenase complex is a multienzyme complex that contains three activities: a pyruvate decarboxylase (E1), a dihydrolipoyl transacetylase (E2), and a dihydrolipoyl dehydrogenase (E3).

- 25 Other α -ketoacid dehydrogenase complexes exist that use the same catalytic scheme with α -ketoacids other than pyruvate. The TCA cycle α -ketoglutarate dehydrogenase complex is an example. The branched-chain α -ketoacid dehydrogenases are also multienzyme complexes

constructed in a manner similar to that of the pyruvate dehydrogenase complex. If the endogenous pyruvate dehydrogenase complex is not sufficiently active in catalyzing the turnover of α -ketobutyrate to propionyl-CoA for copolymer production, it can be modified to enhance its activity.

One way to accomplish this, for example, is to overexpress a branched-chain α -ketoacid decarboxylase E1 subunit having better binding and decarboxylating properties with α -ketobutyrate than does the native PDC E1. For example, the branched chain α -ketoacid dehydrogenase complex of bovine kidney has substantial activity with α -ketobutyrate as substrate, comparable to that with the normal substrate α -ketoisovalerate ($K_m = 56 \mu\text{M}$ and $40 \mu\text{M}$, respectively; specific activity = $12 \mu\text{mol}/\text{min}\cdot\text{mg}$, Pettit et al., 1978). Other examples of branched chain α -ketoacid dehydrogenase complexes are those from *Pseudomonas putida* (Burns et al, 1988), and *Bacillus subtilis* (Lowe et al, 1983). An overexpressed branched-chain α -ketoacid decarboxylase E1 subunit could effectively compete with the endogenous pyruvate dehydrogenase complex E1 subunit, and combine with the pyruvate dehydrogenase complex E2E3 subcomplex to create a functional hybrid complex. This has in fact been shown to occur naturally with the *E. coli* α -ketoglutarate dehydrogenase complex, where approximately 10% of the total E1 component is the pyruvate dehydrogenase complex E1 (Steginsky et al., 1985). The pyruvate dehydrogenase complex E1 component accounts for the pyruvate activity of the α -ketoglutarate dehydrogenase complex. An artificially produced hybrid of E1 (pyruvate decarboxylase) with E2E3 (dihydrolipoyl transsuccinylase-dihydrolipoyl dehydrogenase subcomplex) has higher turnover rates with pyruvate compared to the naturally isolated α -

ketoglutarate dehydrogenase complex, but understandably lower rates than that of the pyruvate dehydrogenase complex itself (Steginsky et al., 1985).

5 Alternatively, both the E1 and E2 components of a branched-chain α -ketoacid dehydrogenase complex that has significant activity with α -keto-butyrate, such as the bovine kidney complex mentioned above, can be expressed. Since the dihydrolipoyl dehydrogenase (E3) is common to all α -ketoacid dehydrogenase complexes, and would be naturally produced for
10 the pyruvate dehydrogenase complex, overexpression of this component may not be necessary. The endproduct would be a fully functional, branched-chain α -keto-acid dehydrogenase complex. If additional E3 is required to interact with overexpressed and endogenous E1 and E2 components, one could overexpress a plant or bacterial E3 (Camp &
15 Randall, 1985, and Camp et al., 1988), or the specific E3 of the branched-chain α -ketoacid dehydrogenase complex, for example the bovine kidney complex (Pettit et al., 1978).

 In addition to an E1 α -ketoacid decarboxylase catalyzing the
20 decarboxylation of an α -ketoacid in an intact complex, unassociated E1 in the presence of an oxidizing agent will catalyze the decarboxylation of an α -ketoacid to form CO_2 and the free acid (Gruys et al., 1989; Speckhard & Frey, 1975; Reed & Willms, 1966). Thus, it is possible that an unassociated branched-chain E1, such as that from bovine kidney, when
25 overexpressed in plants or bacteria, will produce propionate and CO_2 from α -ketobutyrate in the presence of an endogenous oxidizing agent such as ferredoxin. Propionate can be activated to propionyl-CoA by endogenous or overexpressed acyl-CoA synthetase (refer to Example 7).

Mutagenesis of the endogenous E1 of the host PDC is another approach to enhancing its activity toward α -ketobutyrate. Recent reports of mutagenesis of the E2 component from *Bacillus* (Wallis & Perham, 1994), *Azotobacter* (Schulze et al., 1992 & 1993), and the human E3 component (Kim and Patel, 1992), have demonstrated that one can modify both the activity of these enzymes towards substrates as well as their binding affinity to the complex. Similar manipulations can be carried out on the E1 component to enhance its catalytic turnover of α -ketobutyrate.

10

Example 7

Increased Propionyl-CoA Production From α -Ketobutyrate Via Overexpression of Pyruvate Oxidase and Acyl-CoA Synthetase

Another method of enhancing the level of propionyl-CoA in bacteria and plants involves, in a first step, the oxidative decarboxylation of α -ketobutyrate (the amount of which can be increased *in vivo* by utilizing a deregulated threonine deaminase as described above in Examples 2-4), to propionate and CO₂, catalyzed by pyruvate oxidase (E.C. 1.2.3.3) (Figure 3). Unpublished results have shown that pyruvate oxidase efficiently catalyzes this reaction (J.E. Cronan, University of Illinois, personal communication). The *poxB* gene of *E. coli* encodes the pyruvate oxidase enzyme (Grabau and Cronan, 1986). Another example is the pyruvate oxidase from *Lactobacillus plantarum* (Miller and Schulz 1993). This homotetrameric protein of 62 kDa has been thoroughly characterized (Chang & Cronan, 1995, and references cited therein).

In a second step following pyruvate oxidase, endogenous bacterial or plastid acyl-CoA synthetases such as acetyl-CoA synthetase will activate the free propionate to propionyl-CoA (Doi et al., 1986; Nakamura et al., 1991). Yeast acetyl-CoA synthetase has been shown to catalyze the

activation of propionate to propionyl-CoA (Patel and Walt, 1987). In addition to yeast, the *A. eutrophus acoE* gene has been cloned and shown to be active with propionyl-CoA (Priefert and Steinbüchel, 1992). If the endogenous acetyl-CoA synthetase activity proves to be insufficient for
5 producing propionyl-CoA, the enzyme, for example that from yeast, can be overexpressed in the plant, or in a bacterial host.

Another system for activating propionate to propionyl-CoA has been described by Rhie and Dennis (1995). In *E. coli*, the actions of acetate
10 kinase (*ackA*) and phosphotransacetylase (*pta*) are predominately responsible for the conversion of propionate to propionyl-CoA. Thus acetyl-CoA synthetase can be replaced by acetate kinase (*ackA*) and phosphotransacetylase (*pta*) in the present invention.

15 Enhanced propionyl-CoA production can thus be achieved by overexpressing any of the foregoing enzymes in a plant or bacterial host. Overexpression of the foregoing enzymes in combination with an overexpressed deregulated threonine deaminase and overexpressed PHA β -ketothiolase, β -ketoacyl-CoA reductase, and PHA synthase enzymes is
20 expected to result in the production of P(3HB-co-3HV) copolymer.

Example 8

Optimization of β -Ketothiolase, β -Ketoacyl-CoA Reductase, and PHA Synthase Activity for Production of 25 P(3HB-co-3HV) Copolymer in Bacteria and Plants

Alcaligenes eutrophus can produce both PHB homopolymer and, when provided with an appropriate precursor such as propionate, P(3HB-co-3HV) copolymer (U.S. Patent No. 4,477,654). It was expected that
30 P(3HB-co-3HV) copolymer would also be produced if *E. coli* transformed

with the *A. eutrophus* PHB biosynthetic genes was provided with the appropriate precursor, for example, propionate or α -ketobutyrate (see Figure 3).

5 The experiment reported in Table 3, *supra*, was designed in part to test this hypothesis. Introduction of a deregulated threonine deaminase (L481F) into *E. coli* was expected to provide α -ketobutyrate for incorporation of a C5 monomer into the PHA polymer. However, the data in Table 3 demonstrate that while the α -ketobutyrate concentration in cells
10 containing the overexpressed threonine deaminase was dramatically elevated, PHA polymer content and composition were essentially identical in both wild-type cells and cells expressing the introduced threonine deaminase. These results suggest the presence of a metabolic block in the conversion of α -ketobutyrate to P(3HB-co-3HV) copolymer.

15 As shown in Figure 3, there are four reactions required to convert α -ketobutyrate to P(3HB-co-3HV) copolymer. These reactions are catalyzed successively by the pyruvate dehydrogenase complex, β -ketothiolase, acetoacetyl-CoA reductase, and PHB synthase. The
20 metabolic block may therefore involve the inability of one of these enzymes to utilize substrate derived from α -ketobutyrate. However, the literature suggests that all of these enzymes possess the appropriate substrate specificities (Bisswanger, 1981; Haywood et al., 1988a; Haywood et al, 1988b; Haywood et al, 1989). Thus, the substrate specificities of some of
25 these enzymes were reevaluated.

**Substrate Specificity of PhbB Acetoacetyl-CoA Reductase from
A. eutrophus and β -Ketothiolases from Various Sources**

PhbB Acetoacetyl-CoA Reductase

5

Analysis of *A. eutrophus* PhbB acetoacetyl-CoA reductase substrate specificity in the forward reaction was carried out using β -ketobutyryl-CoA (acetoacetyl-CoA) and β -ketovaleryl-CoA. These β -ketoacyl-CoAs were reduced to the corresponding D- β -hydroxyacyl-CoAs using NADPH (Figure 1). This analysis was previously performed by Haywood et al. (1988a), but in the reverse reaction where *in situ*-generated D- β -hydroxyacyl-CoA was oxidized with NADP⁺. It is important to establish that this enzyme can catalyze the forward reaction with β -ketovaleryl-CoA and NADPH to be confident that it does not represent a metabolic block in copolymer production.

E. coli transformed with plasmid pMON25628 (Figure 20) encoding *A. eutrophus* PhbB acetoacetyl-CoA reductase were grown on LB to early log phase and induced with nalidixic acid (60 μ g/ml). Protein extracts were obtained by pelleting induced cells, resuspending in 50mM KPi and 5% glycerol, sonicating, and removing cellular debris by centrifugation. The assay was conducted at pH 7.0 with 100 mM KPi, 0.15 mM NADPH, and 60 μ M β -ketoacyl-CoA. The reaction was initiated with acetoacetyl-CoA reductase. Synthesis of β -ketovaleryl-CoA was accomplished through a scaled-up version (10 μ mol of acyl-CoA in 2 ml total volume) of the *in situ*-generated β -ketoacyl-CoA procedure described below in the section entitled "Thiolysis Activity of *A. eutrophus* BktB." Purification was accomplished through semi-prep C8-reverse-phase HPLC using the following gradient per sample run (Buffer A is 100 mM ammonium acetate, pH 6.0, B is

acetonitrile) with a 4.0 ml/min flow rate: 0-25 min, 5-45% B; 25-30 min, hold at 45% B; 30-32 min, 45-5% B; 32-42 min, hold at 5% B. β -ketovaleryl-CoA eluted and was collected from 8.5 to 10.5 min.

5 Rate results for β -ketobutyryl-CoA and β -ketovaleryl-CoA with PhbB acetoacetyl-CoA reductase gave specific activities of 103 and 28 units/mg protein, respectively, which is 27% relative rate for the C5 versus the C4 substrate under the conditions of the assay. This is consistent with the results described by Haywood et al. (1988a), but further demonstrates
10 that this enzyme is catalytically sufficient for the metabolic conversion of β -ketovaleryl-CoA to D- β -hydroxyvaleryl-CoA in the production of copolymer.

15 **β -Ketothiolases**

Condensation Activity of *A. eutrophus* PhbA and BktB

20 The data presented above demonstrate that the substrate specificity of PhbB is sufficient for production of C5 monomer for incorporation into PHA. Therefore, the substrate specificity of PhbA, as well as of other *A. eutrophus* β -ketothiolases, was investigated.

25 As noted above, *A. eutrophus* can produce P(3HB-co-3HV) copolymer when provided with an appropriate C5 precursor. Slater et al. (1988) demonstrated the presence of genes other than *phbA* encoding β -ketothiolase activity in *A. eutrophus*. One of these β -ketothiolases, designated BktB herein, was encoded on plasmid pBK6. *A. eutrophus* BktB
30 was obtained from *E. coli* DH5 α transformed with plasmid pMON25754. pMON25754 was produced from the *A. eutrophus* pBK6 clone (Slater et al.,

1988) by deleting a 5.0kb XhoI fragment (there is an additional XhoI site approximately 5.0kb 5' to the XhoI site shown in Figure 1 of Slater et al.). The pBK6 clone was produced by Slater et al. (1988) from *A. eutrophus* H16, i.e., ATCC accession number 17699, which is publicly available from the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852. BktB was overexpressed from its native promoter on the high copy plasmid pMON25754 by growing the cells overnight at 37°C to stationary phase in LB broth containing ampicillin (100µg/ml). *A. eutrophus* PhbA was obtained from *E. coli* DH5α transformed with pMON25636 (Figure 22) by growing the cells at 37°C in LB broth containing ampicillin (100µg/ml) to early log phase and then inducing with IPTG. Cells were harvested after two hours of further growth. PhbA and BktB protein extracts were prepared by pelleting the cells, resuspending in 50mM KPi and 5% glycerol, sonicating, and removing cellular debris by centrifugation.

The activity of β-ketothiolases can be assayed in both the condensation and thiolysis directions. However, condensation activity for products that are the result of a mixed condensation of acetyl-CoA and a longer chain acyl-CoA are difficult to measure. This is due to the difficulty in distinguishing between acetoacetyl-CoA, the product of condensing two acetyl-CoA molecules, and the mixed condensation reaction product, both of which form simultaneously. To date, condensation assays have been spectrophotometrically-based, but these are valid only when there is a single acyl-CoA substrate present (i.e., acetyl-CoA) since they cannot discriminate between the products formed. The assay described below represents the first reported assay that permits complete quantitation of all condensation products regardless of the nature of starting substrates.

β -ketothiolase assays designed to measure the condensation activity with acetyl-CoA plus either propionyl-CoA or butyryl-CoA were performed using 1- 14 C-labelled acetyl-CoA. Since the equilibrium for the reaction lies heavily toward the starting acyl-CoAs rather than the condensed β -ketoacyl-CoA product, the latter was "pulled" to the corresponding β -hydroxyacyl-CoA using NADH and β -ketoacyl-CoA dehydrogenase. In addition, to avoid the strong feedback inhibition of β -ketothiolase by free CoA, each assay solution contained 5,5'-dithio-bis(2-nitrobenzoic acid) (DTNB) to trap chemically the free CoA produced by the reaction.

The condensation assay mixture (0.5 ml final volume) contained 100 mM KPi, pH 7.8, 0.5 mM NADH, 0.30 mM 1- 14 C-acetyl-CoA (0.15 μ Ci), 5 units β -ketoacyl-CoA dehydrogenase, 0.5 mM DTNB, and variable propionyl- or butyryl-CoA as indicated in Table 5. The reaction was initiated by the addition of β -ketothiolase. Following 20 minutes at 30°C, each reaction mixture was quenched by the addition of 25 μ l of 10% formic acid. Thirty μ l of 25% H_2SO_4 were added to 200 μ l of quenched reaction mix; this mixture was then incubated at 70°C for 24 hours to hydrolyze all CoA thioesters to their free acids. Sixty μ l of 5 M sodium formate were added to the sample to neutralize the pH, followed by analysis of a 130 μ l aliquot using C18-reverse-phase HPLC (0.5 x 25 cm column). Detection and quantitation were carried out by monitoring flow-through with a Radiomatic flow detector. A gradient elution was employed using aqueous 1% acetic acid (solvent A) and acetonitrile (solvent B). The following gradient program was used per sample run with a flow rate of 1.0 ml/min: 0-15 min, 0-35% B; 15-20 min hold at 35% B; 20-21 min, 35-0% B; 21-30 min hold at 0% B. Retention times for 14 C-labelled compounds are as follows; acetic acid, 5.1 min; β -hydroxybutyric acid, 7.5 min; β -

hydroxyvaleric acid, 11.7 min; and β -hydroxycaproic acid, 15.3 min.

Calculation of the percent turnover of 1-¹⁴C-acetyl-CoA to product was based on the ¹⁴C-peak integration, correcting for the double label which occurs in β-hydroxybutyric acid (i.e., from two molecules of

5 1-¹⁴C-acetyl-CoA). The results are shown in Table 5.

[illegible]

TABLE 5
Substrate Specificity of *A. eutrophus* PhbA and BktB
 β -Ketothiolases in the Condensation Reaction
with Acetyl-CoA plus Propionyl-CoA or Butyryl-CoA¹

5

| Enzyme Sample | Propionyl- CoA (mM) | % Turnover C4 Product | % Turnover C5 Product | % Turnover C6 Product |
|------------------|------------------------------|--------------------------|--------------------------|--------------------------|
| PhbA | 0 | 16.9 | | |
| PhbA | 0.6 | 13.5 | 0 | |
| PhbA | 1.2 | 10.3 | 0 | |
| PhbA | 2.4 | 7.0 | 0 | |
| BktB | 0 | 4.2 | | |
| BktB | 0.6 | 1.8 | 13.2 | |
| BktB | 1.2 | 1.0 | 15.6 | |
| BktB | 2.4 | 0.7 | 17.8 | |
| | Butyryl- CoA (mM) | | | |
| BktB | 0.6 | 1.0 | | 4.9 |
| BktB | 1.2 | 0.6 | | 7.3 |
| BktB | 2.4 | 0.4 | | 7.9 |

¹ For all reactions, the acetyl-CoA concentration was 0.30 mM (0.15 μ Ci ¹⁴C-acetyl-CoA per 0.50 ml reaction). The concentration of propionyl-CoA or butyryl-CoA refers to the concentration in the assay. % turnover to product refers to the amount of acetyl-CoA converted to C4, C5, or C6 product. PhbA β -ketothiolase was not tested for C6 product formation with butyryl-CoA. Blank spaces indicate products that were not detected in the experiment, probably because the appropriate acyl-CoAs were not present in the individual reaction mixture.

10

The data in Table 5 clearly demonstrate that the β -ketothiolase activity of PhbA is restricted to the catalytic formation of the C4 compound acetoacetyl-CoA. These results can explain the metabolic block to P(3HB-co-3HV) copolymer production in recombinant *E. coli* containing *phbA*, *phbB*, *phbC* and *E. coli* *ilvA466* (L481F) (Table 3). The data in Table 5 for *A. eutrophus* PhbA contrast with the suggestions of Haywood et al. (1988b) that propose, based on thiolysis data, that the activity of PhbA β -ketothiolase alone can account for the β -ketovaleryl-CoA needed for the C5 component in P(3HB-co-3HV) copolymer production. The data presented herein demonstrate that PhbA cannot efficiently catalyze the formation of the C5 monomer. Thus, a β -ketothiolase other than PhbA appears to be involved in the production of P(3HB-co-3HV) copolymer in *A. eutrophus*.

As shown in Table 5, the *A. eutrophus* BktB enzyme catalyzes the production of C4, C5, and C6 β -ketoacyl-CoA compounds, with an apparent preference for the formation of the C5 product. These results suggest that the biochemical pathway of P(3HB-co-3HV) copolymer production in *A. eutrophus* involves the BktB β -ketothiolase and/or another β -ketothiolase capable of forming the C5 product. These results further suggest that the biosynthesis of significant amounts of P(3HB-co-3HV) copolymer in heterologous host bacteria and plants can be facilitated by the use of a β -ketothiolase having condensation substrate specificity similar or identical to that of BktB, i.e., being capable of condensing acetyl-CoA with propionyl-CoA to form β -ketovaleryl-CoA.

25

In addition to its ability to catalyze the formation of β -ketovaleryl-CoA, the BktB β -ketothiolase can also catalyze the formation of β -keto-caproyl-CoA (Table 5). Based on this observation, the BktB β -ketothiolase

can also be utilized for the production of copolymers containing monomers ranging in size from C4 to C6. Although the condensing activity of the BktB β -ketothiolase beyond C6 has not been tested, it is possible that this enzyme can catalyze the formation of β -ketoacyl-CoA compounds greater
5 than C6. It is likely that plant and bacterial cells containing BktB and also expressing appropriate β -ketoacyl-CoA reductases and PHA synthases capable of utilizing the range of β -keto-acyl-CoA substrates produced by BktB (for example, *Nocardia corallina* PHA synthase; Dennis, 1994) can produce copolymers containing monomer units ranging from C4 to C6, and
10 possibly beyond. One example is a copolymer comprising β -hydroxybutyrate (3HB) and β -hydroxycaproate (3HC), designated P(3HB-co-3HC) copolymer.

Thiolysis Activity of *A. eutrophus* BktB

15

To enhance further our understanding of the kinetic characteristics of BktB β -ketothiolase, a thorough kinetic analysis of the enzyme's thiolysis activity was conducted using purified protein. The assay to determine the kinetic parameters K_m and V_{max} for β -ketobutyryl-CoA (C4),
20 β -ketovaleryl-CoA (C5), and β -ketocaproyl-CoA (C6), was performed at 25°C by the *in situ* generation of the β -keto compounds from their corresponding trans-2,3-enoyl-CoAs essentially as described by Haywood et al. (1988). Utilization of variable substrate concentrations permitted the determination of the descriptive kinetic parameters of the β -
25 ketothiolase. Kinetic rates were determined only at fixed substrate concentrations in the Haywood et al. communication.

Purified BktB protein was prepared by growing *E. coli* EE36 transformed with pMON25754 in LB broth containing ampicillin (100µg/ml) overnight at 37°C. Cells were then harvested by centrifugation, and the pellet was resuspended in 50mM KPi and 5% glycerol, sonicated, and the sonicate centrifuged to remove cellular debris. The cell extract was desalted using a PD-10 column (Pharmacia), and proteins resolved using a Pharmacia FPLC system and MonoQ anion-exchange column (Pharmacia). Buffer A was 10 mM Tris buffer, pH 7.8, plus 1mM DTT; buffer B was buffer A plus 1 M KCl. The gradient was 0 to 10% B in 10 min, then to 35% B in 40 min. The flow rate was 1 ml/min, and the resolved proteins were collected at the rate of 1 ml/fraction. The BktB protein was found in fractions 20 to 27, with the maximum activity peak at fraction 23. Fraction 23 used for N-terminal sequencing was desalted, washed, and stored in 10 mM sodium bicarbonate using a Centricon-10 concentrator (Amicon, Inc.) Protein was found to be pure according to SDS-PAGE analysis, with an apparent molecular weight of 39.5 kD (data not shown). The enzyme was stored in 10mM Tris-HCl, pH 7.8, 1mM DTT, 50% glycerol at -20 °C. Activity remained constant for a minimum of two months under these conditions.

20

The purified BktB β-ketothiolase was diluted to approximately 1 mg/ml in 10mM Tris-HCl, pH 7.8, 50% glycerol. Two rabbits were boosted initially with 200 µg of purified protein in a total volume of 1 ml by Scientific Associates Inc. (St. Louis, MO). Bleeds after the first boost yielded polyclonal antibodies which produced a single immunoreactive band at approximately 45 kD on Western blots containing a range of dilutions of purified BktB. The detection limit of the antibodies was determined to be 1 ng. These antibodies can be used to detect proteins having similar immunoreactive properties to BktB β-ketothiolase, thus making it possible to detect ketothiolases having similar enzymatic activity to BktB in crude extracts.

The C5 and C6 enoyl-CoAs were synthesized according to Schulz (1974) and purified using semi-prep C8-reverse-phase HPLC as described above for β -ketovaleryl-CoA. β -ketobutyryl-CoA (acetoacetyl-CoA) was purchased from Sigma (St. Louis, MO). The assay mixture (1 ml final volume) contained 150 mM EPPS, pH 8.0, 50 mM MgCl_2 , 1.5 mM pyruvate, 0.4 mM NAD^+ , and a coupling enzyme cocktail of three units crotonase, five units β -hydroxyacyl-CoA dehydrogenase, and 10 units lactic dehydrogenase. The enoyl-CoA was added to this mixture and the absorbance at 304 nm was monitored until it reached a plateau (typically about 5 min). CoA was added to this mixture, followed by BktB β -ketothiolase to initiate the reaction. The decrease in absorbance at 304 nm due to the thiolysis of the β -ketoacyl-CoA was utilized as a direct measure of activity. Extinction coefficients to convert absorbance units to μmol of product were 19.5, 12.2, and 14.0 $\text{cm}^{-1} \text{mM}^{-1}$ for β -ketobutyryl-CoA, β -ketovaleryl-CoA, and β -ketocaproyl-CoA, respectively. β -ketoacyl-CoA concentration was varied from 5 to 80 μM (CoA was fixed at 200 μM), and CoA concentration was varied from 20 to 400 μM (β -ketocaproyl-CoA was fixed at 60 μM) for determining kinetic parameters for β -ketoacyl-CoA and CoA, respectively. Reaction rates were calculated in the steady-state within the first minute following initiation. All data were fitted to the normal Michaelis-Menten equation using a non-linear regression analysis. The results are shown in Table 6.

TABLE 6
Kinetic Parameters for BktB β -Ketothiolase
in the Thiolytic Reaction

| Substrate | K_m (μ M) | V_{max} (u/mg) ¹ | V_{max} / K_m |
|--------------------------------|------------------|-------------------------------|-----------------|
| B-ketobutyryl-CoA ² | 37 ± 3 | 122 ± 5 | 3.0 |
| B-ketovaleryl-CoA ² | 15 ± 1 | 236 ± 8 | 16 |
| B-ketocaproyl-CoA ² | 6.0 ± 0.7 | 103 ± 3 | 17 |
| CoA ³ | 53 ± 6 | 98 ± 3 | 1.8 |

¹ A unit, u, refers to μ mol product formed per minute.

² Kinetic parameters were determined using a fixed concentration of CoA of 200 μ M.

³ Kinetic parameters were determined using a fixed concentration of β -ketocaproyl-CoA of 60 μ M.

The results in Table 6 show that β -ketovaleryl-CoA can be turned over by BktB by about a factor of two faster than the other two substrates according to the V_{max} values. This faster turnover is somewhat modulated by a higher K_m compared to β -ketocaproyl-CoA. Overall, based on V_{max} / K_m , the enzyme utilizes C5 and C6 substrates about equivalently, whereas β -keto-butyryl-CoA is approximately five-fold less efficient in catalytic turnover. These results are consistent with the condensation data for BktB shown in Table 5. From this, and in combination with the condensation and thiolysis results on PhbA β -ketothiolase, one might hypothesize that gathering thiolysis data alone for a particular β -ketothiolase would be sufficient to suggest the activity of the enzyme in the condensation direction. However, as shown below, this is not necessarily true.

Thiolysis and Condensation Activity of Other β -Ketothiolases

Two additional β -ketothiolases from *A. eutrophus* (cosmid clones AE65 and AE902; Slater et. al., 1988), plus two β -ketothiolases from

- 5 *Zoogloea ramigera*, were analyzed for thiolysis and condensation activity using the assays described above. *Zoogloea ramigera* (ATCC 19623) was grown at 30°C for 48 hours in medium containing; 9.4 g/l K_2HPO_4 ; 2.2 g/l KH_2PO_4 ; 2.5g/l $MgSO_4 \cdot 7H_2O$; 3g/l N-Z Amine; and 10g/l Na-gluconate. *E. coli* containing cosmid clones AE65 and AE902 were grown for 16 hours at
- 10 37°C in LB supplemented with 50 μ g/ml kanamycin. For the thiolysis assay, activity was monitored at a fixed concentration of β -ketoacyl-CoA at 40 μ M and CoA at 100 μ M. All of the protein extracts were partially purified using the Pharmacia FPLC MonoQ separation described above. Table 7 shows the thiolysis results.

15

TABLE 7
Substrate Specificity of Various β -Ketothiolases
in the Thiolysis Reaction¹

20

| Enzyme Sample | B-ketobutyryl-CoA (u/mg) | B-ketovaleryl-CoA (u/mg) | B-ketocaproyl-CoA (u/mg) |
|-------------------------------------|--------------------------|--------------------------|--------------------------|
| <i>Z. ramigera</i> (A) ² | 0.41 | 0.15 | 0.02 |
| <i>Z. ramigera</i> (B) ² | 0.21 | 0.19 | 0.11 |
| pAE65 | 0.156 | 0.502 | 3.16 |
| pAE902 | 0.585 | 1.25 | 0.561 |

¹ Activity was determined using partially purified enzymes. The concentration of substrate was 40 μ M. pAE designations are *A. eutrophus* cosmid clones.

- 25 ² The *Z. ramigera* protein sample yielded two active ketothiolase peaks upon FPLC MonoQ resolution, designated A and B.

- MonoQ separation of the *Z. ramigera* protein extract resulted in two resolved active β -ketothiolase peaks. The first peak, designated A, shows thiolysis activity with C4 and C5 substrate, but little if any with C6. Based on this activity, this β -ketothiolase in all likelihood is the enzyme purified
- 5 and described by Davis et al., (1987), and is the protein expressed specifically for PHB production in this organism. *Z. ramigera* peak B, which has not been previously described, appears to have broadened specificity, at least to C6.
- 10 *A. eutrophus* cosmid clones pAE65 and pAE902, when similarly analyzed for thiolysis activity, both showed specificity at least to C6. Clone pAE65, in fact, likely has significant activity with β -ketoacyl-CoAs of higher chain-length than C6 based on the significant increase in activity when going from C4 to C6. Additional results (not shown) for pAE65
- 15 demonstrate dehydratase and β -keto-acyl-CoA dehydrogenase activity in the same resolved MonoQ fraction. This suggests that the pAE65 clone contains the genes for the fatty acid β -oxidative complex proteins of *A. eutrophus*. Interestingly, clone pAE902 encodes an enzyme having thiolysis activity with β -ketoacyl-CoA substrates ranging from C4 to C6
- 20 similar to that of BktB. As such, considering only the tholysis results in Table 7, pAE902 might be considered a good candidate for β -ketovaleryl-CoA condensation production from acetyl-CoA and propionyl-CoA. However, it is important to consider condensation direction activity as well. Condensation results for this clone, along with that of the other β -
- 25 ketothiolases, are shown in Table 8.

TABLE 8
Substrate Specificity of Various β -Ketothiolases
in the Condensation Reaction with
Acetyl-CoA Plus Propionyl-CoA or Butyryl-CoA

5

| Enzyme Sample | Substrates ¹ | % Turnover C4 Product | % Turnover C5 Product | % Turnover C6 Product |
|-------------------------------------|-------------------------|-----------------------|-----------------------|-----------------------|
| <i>Z. ramigera</i> (A) ² | C2 | 23.3 | | |
| <i>Z. ramigera</i> (A) ² | C2 + C3 | 12.4 | 7.1 | |
| <i>Z. ramigera</i> (A) ² | C2 + C4 | 18.9 | | 0 |
| <i>Z. ramigera</i> (B) ² | C2 | 3.4 | | |
| <i>Z. ramigera</i> (B) ² | C2 + C3 | 1.1 | 2.7 | |
| <i>Z. ramigera</i> (B) ² | C2 + C4 | 1.2 | | 2.6 |
| pAE65 | C2 | 5.7 | | |
| pAE65 | C2 + C3 | 2.2 | 58.6 | |
| pAE65 | C2 + C4 | 0.5 | | 84.3 |
| pAE902 | C2 | 0 | | |
| pAE902 | C2 + C3 | 0 | 0 | |
| pAE902 | C2 + C4 | 0 | | 0 |

¹ For all reactions, the acetyl-CoA (C2) concentration was 0.40 mM (0.15 μ Ci ¹⁴C-acetyl-CoA per 0.50 ml reaction). The concentration of propionyl-CoA (C3) or butyryl-CoA (C4) was 2.0 mM when present. % turnover to product refers to the amount of acetyl-CoA converted to C4, C5, or C6 product. pAE designations are *A. eutrophus* cosmid clones.

² The *Z. ramigera* protein sample yielded two active ketothiolase peaks upon FPLC Mono-Q resolution, designated A and B.

15 The data presented in Table 8 demonstrate condensation activity for all samples tested, excluding pAE902. Consistent with the pattern shown in the thiolysis results, the *Z. ramigera* enzymes are capable of producing C5, and C5 and C6 condensation products, for peaks A and B, respectively. These then represent two possible alternative candidates for
20 use in the biosynthesis of P(3HB-co-3HV) copolymer in recombinant

systems. Similarly, cosmid clone pAE65 shows activity that parallels that seen in the thiolysis assay.

The pAE902 cosmid clone showed no condensation products, and in addition exhibited greater than 50% hydrolysis of the starting ^{14}C -acetyl-CoA in the time-course of the reaction (hydrolysis data not shown). This result shows that it is important to demonstrate condensation substrate specificity activity for a particular β -ketothiolase to evaluate whether or not it is useful in producing P(3HB-Co-3HV) copolymer. That is, from this result with pAE902, it is not obvious that the thiolysis activity of a particular β -ketothiolase will necessarily translate to useful information regarding its condensation activity. This also demonstrates the utility of a quantitative assay such as the one described herein for measuring condensation activity.

Cloning of the *A. eutrophus* *bktB* Gene

As noted above, Slater et. al. (1988) identified several putative *A. eutrophus* β -ketothiolase clones, including pBK6, by screening *E. coli* harboring an *A. eutrophus* cosmid library for β -ketothiolase activity.

The β -ketothiolase gene encoded on pBK6 was mapped by serial subcloning and deletion analysis, and sequenced (Sambrook et al., 1989). The sequence is shown in SEQ ID NO:9. For the purpose of the present invention, this gene has been designated as the *bktB* gene of *A. eutrophus*. This gene was deposited under the terms of the Budapest Treaty in the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852, U.S.A., as an insert in plasmid pMON25728 contained in *E. coli* DH5 α on March 6, 1996, designated ATCC 98007.

pMON25728 contains a fragment of approximately 3.9 Kb extending from a Bgl II site approximately 550 bp upstream of the *bktB* open reading frame to a Nco I site downstream of *bktB*. There is a single additional Nco I site in this fragment, and it is within *bktB*. The Bgl II site upstream of *bktB* is the single Bgl II site shown in plasmid pBK6 (Slater et al, 1988). pMON25728 contains the 3.9 Kb Bgl II-NcoI fragment in which both sites have been end-filled and ligated to DNA of pBluescript KS+ (Stratagene). pBluescript KS+ polylinker sites were cut with Sac I and Eco RV, and the Sac I site was treated with exonuclease to create a blunt end. The Bgl II site of the *A. eutrophus* DNA has been fused to the EcoRV site of pBluescript KS+.

Automated Edman degradation chemistry was used to determine the NH₂-terminal protein sequence (Hunkapiller et al., 1983) of purified BktB protein. A Perkin Elmer Applied Biosystems Division sequencer (Foster City, CA) was employed for the analysis. The respective PTH-aa derivatives were identified by RP-HPLC analysis in an on-line fashion employing a Brownlee 2.1 mm I.D. PTH-C18 column.

Amino acid sequencing of the protein purified (0.9 mg/ml) as described above revealed that the N-terminal sequence minus the initiating methionine (TREVVVVSGVRTAIG, SEQ ID NO:10) corresponds to that predicted by the DNA sequence shown in SEQ ID NO:9, as well as the deduced amino acid sequence shown in SEQ ID NO:11.

P(3HB-co-3HV) Copolymer Production and Active Expression of *bktB* in *E. coli*

The data reported in Table 3 (Example 2) suggested the presence of a metabolic block to P(3HB-co-3HV) copolymer production in *E. coli*

transformed with the *A. eutrophus phbA*, *phbB*, *phbC*, and *E. coli ilvA466* (L481F) genes. To determine whether the expression of the *A. eutrophus bktB* gene in *E. coli* is catalytically effective in P(3HB-co-3HV) copolymer biosynthesis *in vivo*, the following experiment was performed.

5

Two sets of *E. coli* DH5 α cells were transformed with the plasmid combinations shown in Table 9 as described in Example 2 in the section entitled "*In vivo* Analysis of Cloned *E. coli ilvA* genes":

10

TABLE 9
Plasmid Combinations Used for the Production
of P(3HB-co-3HV) Copolymer in Transformed *E. coli*

| Set 1 | Gene | Antibiotic resistance | Promoter | Origin of replication |
|--------------|-------------|-----------------------|----------|-----------------------|
| pMON25628 | <i>phbB</i> | Spectinomycin | recA | ori327 |
| pMON25629 | <i>phbC</i> | Kanamycin | tac | ori pACYC |
| pMON25636 | <i>phbA</i> | Ampicillin | tac | ori CoLE1 |
| Set 2 | | | | |
| pMON25628 | <i>phbB</i> | Spectinomycin | recA | ori327 |
| pMON25629 | <i>phbC</i> | Kanamycin | tac | ori pACYC |
| pMON25728 | <i>bktB</i> | Ampicillin | native | ori CoLE1 |

15

Maps of pMON25628, pMON25629, and pMON25636 are shown in Figures 20-22, respectively. pMON25728, a derivative of pBK6 (Slater

et al., 1988), was created by first subcloning a 5.5Kb NotI/EcoRI fragment of pBK6 into pBluescript KS+ (Stratagene), creating pMON25722.

pMON25722 was subsequently digested with BglII and EcoRI, releasing a 1.1Kb fragment. The remaining vector was blunt ended with Klenow and

- 5 religated, creating pMON25724. Finally, pMON25724 was partially digested with NcoI, and digested to completion with SacI, releasing a 0.9Kb fragment. The remaining vector was blunt ended with Klenow and religated, creating pMON25728.

- 10 Each set of transformed cells containing all three plasmids was selected by the addition of all three antibiotics (ampicillin, 100 µg/ml; kanamycin, 50 µg/ml; spectinomycin, 75 µg/ml) to the growth medium (LB agar). Single isolated colonies were grown on LB medium containing all three antibiotics to early log phase and induced with IPTG and nalidixic
- 15 acid. Cells were pelleted by centrifugation, resuspended in 500µl of 50mM KPi/5% glycerol, and sonicated on ice. Enzyme assays as described above were performed on the selected cells to confirm expression and activity of the reductase (PhbB), and, in the thiolysis direction, the β-ketothiolases (PhbA and BktB). The results are shown in Table 10.

20

TABLE 10

Activity of PhbA, BktB, and PhbB in Transformed *E. coli* DH5α

| Set 1 | Gene | <u>Thiolase Rate</u> (AU/min ³) C4 substrate ¹ | <u>Thiolase Rate</u> (AU/min ³) C5 substrate ² | <u>Reductase</u> Rate (AU/min ³) C4 substrate ¹ |
|--------------|-------------|---|---|--|
| pMON25628 | <i>phbB</i> | | | 0.035 |
| pMON25629 | <i>phbC</i> | | | |
| pMON25636 | <i>phbA</i> | 0.330 | 0.013 | |
| Set 2 | | | | |

| | | | | |
|-----------------------------|-------------|-------|-------|-------|
| pMON25628 | <i>phbB</i> | | | 0.051 |
| pMON25629 | <i>phbC</i> | | | |
| pMON25728 | <i>bktB</i> | 0.132 | 0.135 | |
| Control | | | | |
| <i>E. coli</i> DH5 α | | 0.008 | | 0.004 |

¹C4 substrate: acetoacetyl-CoA

²C5 substrate: β -ketovaleryl-CoA

³AU refers to absorbance units

5

The activities of the PhbA and BktB enzymes were as expected based upon the kinetic data reported in Table 5, i.e., the PhbA β -ketothiolase was not active in thiolysis of the C5 substrate, while the BktB β -ketothiolase exhibited significant activity with C5 substrate. The levels

10 of reductase (PhbB) activity were low due the nature of the *E. coli* DH5 α host cells. Nalidixic acid is commonly used as a chemical inducer of the SOS response of *E. coli* (Walker, 1987). Since the *phbB* gene is transcribed from the SOS responsive *recA* promoter, and since *E. coli* DH5 α are *recA*, there was no induction from the RecA promoter upon the addition of

15 nalidixic acid. The low levels of reductase activity observed are due to leaky expression from the *recA* promoter. PhbC activity was not measured enzymatically; instead, the production of PHA within the *E. coli* cells was taken as proof of active PhbC enzyme.

20 *E. coli* DH5 α cells containing both sets of plasmids expressing all of the required PHA biosynthesis enzymes were grown on 50ml of LB broth containing all three antibiotics supplemented with 0.75% glucose and 0.15% propionate. The tac promoters were induced with IPTG at a final concentration of 0.5 mM. Nalidixic acid was added to a final concentration

25 of 60 μ g/ml. The cells were grown under antibiotic selection (see above) for 24 hours and assayed for the presence of PHA polymer. Cells were

centrifuged at 7,000 rpm for 20 min., washed with 25 ml methanol, recentrifuged, washed with 15 ml hexane, and recentrifuged. Cell pellets were dried under N₂ for two hours, and the dry cell weights determined. 6.5 ml of chloroform were added to extract the PHA at 100°C for one hour. The solution was cooled and filtered through a PTFE syringe filter (13mm diameter, 0.45µm pore). The PHA was precipitated by the addition of 50 ml of methanol, centrifuged, and washed with hexane. The polymer was dried at 70°C for two hours, and polymer weight was determined. Methanolysis was performed by dissolving 3-5mg of PHA sample by the addition of 1 ml of CHCl₃, with methyl benzoate as an internal standard for normalization of the data. One ml of 15% H₂SO₄/MeOH was added, and the mixture was heated at 100°C for two hours. The samples were cooled, and 0.5 ml of H₂O were added. The lower CHCl₃ layer was removed, and excess water was removed using Na₂SO₄. The solution was then analyzed by gas chromatography as described in the section entitled "*In vivo* Analysis of Cloned *E. coli* *ilvA* Genes". The results are shown in Table 11.

TABLE 11

Gas Chromatographic Analysis of PHA Produced by Transformed *E. coli* DH5α

| Set | Cell weight (mg) | PHA weight (mg) | %PHA | %C4 | %C5 |
|---|------------------|-----------------|------|------|-----|
| Set 1 (<i>phbA</i> + <i>phbB</i> + <i>phbC</i>) | 146.5 | 38 | 26 | 99.1 | 0.9 |
| Set 2 (<i>BktB</i> + <i>phbB</i> + <i>phbC</i>) | 132.4 | 32 | 24 | 95 | 5 |

The results in Table 11 demonstrate that the total amount of P(3HB-co-3HV) copolymer produced in cells transformed with both sets of plasmid vector combinations was comparable. Furthermore, cells expressing the BktB β -ketothiolase produced a greater percentage (five-to-six-fold) of the C5 monomer than cells expressing the PhbA β -ketothiolase. In addition, the formation of P(3HB-co-3HV) in the experiments reported in Table 11 was confirmed by $^1\text{H-NMR}$ (data not shown). These results support the hypothesis that the *A. eutrophus* PhbA β -ketothiolase was responsible for the metabolic block preventing the production of significant C5 constituent of the copolymer reported in Table 3.

Other β -Ketothiolases, β -Ketoacyl-CoA Reductases, and PHA Synthases for the Production of P(3HB-co-3HV) Copolymer

Examples of enzymes useful in the production of P(3HB-co-3HV) copolymer have been described above. However, the present invention is not limited thereto, and other β -ketothiolases, β -ketoacyl-CoA reductases, and PHA synthases that can be used in the present invention exist, or can be obtained from other sources.

One source includes organisms possessing useful PHA biosynthetic enzymes. These include, for example, *Alcaligenes eutrophus*, *Alcaligenes faecalis*, *Aphanothece* sp., *Azotobacter vinelandii*, *Bacillus cereus*, *Bacillus megaterium*, *Beijerinckia indica*, *Derxia gummosa*, *Methylobacterium* sp., *Microcoleus* sp., *Nocardia corallina*, *Pseudomonas cepacia*, *Pseudomonas extorquens*, *Pseudomonas oleovorans*, *Rhodobacter sphaeroides*, *Rhodobacter capsulatus*, *Rhodospirillum rubrum* (Brandl et al., 1990; Doi, 1990), and

Thiocapsa pfennigii. Using the methods described herein, one can identify and isolate DNAs encoding other β -ketothiolases, β -ketoacyl-CoA reductases, and PHA synthases useful in the present invention from these and other organisms capable of producing PHAs.

5

The present invention encompasses not only the *A. eutrophus* DNA sequence shown in SEQ ID NO:9, but also biologically functional equivalent nucleotide sequences. The phrase "biologically functional equivalent nucleotide sequences" denotes DNAs and RNAs, including chromosomal DNA, plasmid DNA, cDNA, synthetic DNA, and mRNA nucleotide sequences, that encode peptides, polypeptides, and proteins exhibiting the same or similar β -ketothiolase enzymatic activity as that of *A. eutrophus* BktB when assayed enzymatically or by complementation. Such biologically functional equivalent nucleotide sequences can encode peptides, polypeptides, and proteins that contain a region or moiety exhibiting sequence similarity to the corresponding region or moiety of the *A. eutrophus* BktB β -ketothiolase.

Thus, one can isolate enzymes useful in the present invention from various organisms based on homology or sequence identity. Tombolini et al. (1995) conducted a comparative study of the homologies of known β -ketothiolases and reductases. Although one embodiment of a nucleotide sequence encoding *A. eutrophus bktB* is shown in SEQ ID NO:9, it should be understood that other biologically functional equivalent forms of *A. eutrophus* BktB-encoding nucleic acids can be readily isolated using conventional DNA-DNA or DNA-RNA hybridization techniques. Thus, the present invention also includes nucleotide sequences that hybridize to SEQ ID NO:9 and its complementary sequence, and that code on expression for peptides, polypeptides, and proteins exhibiting the same or similar enzymatic activity as that of *A. eutrophus* BktB β -ketothiolase. Such nucleotide sequences preferably hybridize to SEQ ID NO:9 or its complementary sequence under moderate to high stringency (see

Sambrook et al., 1989). Exemplary conditions include initial hybridization in 6X SSC, 5X Denhardt's solution, 100 µg/ml fish sperm DNA, 0.1% SDS, at 55°C for sufficient time to permit hybridization (e.g., several hours to overnight), followed by washing two times for 15 min each in 2X SSC, 0.1% SDS, at room temperature, and two times for 15 min each in 0.5-1X SSC, 0.1% SDS, at 55°C, followed by autoradiography. Typically, the nucleic acid molecule is capable of hybridizing when the hybridization mixture is washed at least one time in 0.1X SSC at 55°C, preferably at 60°C, and more preferably at 65°C.

The present invention also encompasses nucleotide sequences that hybridize under salt and temperature conditions equivalent to those described above to genomic DNA, plasmid DNA, cDNA, or synthetic DNA molecules that encode the amino acid sequence of *A. eutrophus* BktB β-ketothiolase, and genetically degenerate forms thereof due to the degeneracy of the genetic code, and that code on expression for a peptide, polypeptide, or protein that has the same or similar ketothiolase enzymatic activity as that of *A. eutrophus* BktB β-ketothiolase.

Biologically functional equivalent nucleotide sequences of the present invention also include nucleotide sequences that encode conservative amino acid changes within the *A. eutrophus* BktB amino acid sequence, producing silent changes therein. Such nucleotide sequences thus contain corresponding base substitutions based upon the genetic code compared to wild-type nucleotide sequences encoding *A. eutrophus* BktB.

In addition to nucleotide sequences encoding conservative amino acid changes within the naturally occurring *A. eutrophus* BktB amino acid sequence, biologically functional equivalent nucleotide sequences of the present invention also include genomic DNA, plasmid DNA, cDNA, synthetic DNA, and mRNA nucleotide sequences encoding non-conservative amino acid substitutions, additions, or deletions. These include nucleic acids that contain the same

inherent genetic information as that contained in the DNA of SEQ ID NO:9, and which encode peptides, polypeptides, or proteins exhibiting the same or similar β -ketothiolase enzymatic activity as that of *A. eutrophus* BktB. Such nucleotide sequences can encode fragments or variants of *A. eutrophus* BktB.

- 5 The *A. eutrophus* BktB β -ketothiolase-like enzymatic activity of such fragments and variants can be identified by complementation or enzymatic assays as described above. These biologically functional equivalent nucleotide sequences can possess from 40% sequence identity, or from 60% sequence identity, or from 80% sequence identity, to 100% sequence identity to naturally occurring DNA or cDNA encoding *A. eutrophus* BktB β -ketothiolase, or corresponding regions or moieties thereof. However, regardless of the percent sequence identity of these biologically functional equivalent nucleotide sequences, the encoded proteins would possess the same or similar enzymatic activity as that of BktB β -ketothiolase. Thus, the biologically functional equivalent nucleotide sequences encompassed by the present invention include sequences having less than 40% sequence identity to SEQ ID NO:9, so long as they encode peptides, polypeptides, or proteins having the same or similar enzymatic activity as that of BktB β -ketothiolase.

- 20 Mutations made in *A. eutrophus* BktB β -ketothiolase cDNA, chromosomal DNA, plasmid DNA, synthetic DNA, mRNA, or other nucleic acid preferably preserve the reading frame of the coding sequence. Furthermore, these mutations preferably do not create complementary regions that could hybridize to produce secondary mRNA structures, such as loops or hairpins, that would adversely affect mRNA translation.

Useful biologically functional equivalent forms of the DNA of SEQ ID NO:9 include DNAs comprising nucleotide sequences that exhibit a level of sequence identity to corresponding regions or moieties of the genomic DNA of

SEQ ID NO:9 from 40% sequence identity, or from 60% sequence identity, or from 80% sequence identity, to 100% sequence identity. However, regardless of the percent sequence identity of these nucleotide sequences, the encoded peptides, polypeptides, or proteins would possess the same or similar

5 enzymatic activity BktB β -ketothiolase. Thus, biologically functional equivalent nucleotide sequences encompassed by the present invention include sequences having less than 40% sequence identity to SEQ ID NO:9, so long as they encode peptides, polypeptides, or proteins having the same or similar enzymatic activity as BktB β -ketothiolase. Sequence identity can be
10 determined, for example, using the "BestFit," "Gap," or "FASTA" programs of the Sequence Analysis Software Package, Genetics Computer Group, Inc., University of Wisconsin Biotechnology Center, Madison, WI 53711, or using the "BLAST" program (Altschul et al., 1990).

15 Useful biologically functional equivalent forms of the BktB protein sequence shown in SEQ ID NO:11 include polypeptides and proteins comprising amino acid sequences that exhibit a level of sequence identity to the BktB protein or regions thereof of at least about 25%, preferably at least about 30%, and more preferably at least about 35%. For this purpose, sequence identity
20 can be determined using the "TFASTA" program of the Sequence Analysis Software Package referred to above, or by the "BLASTP" program of Altschul et al., 1990, for example. Useful biologically functional equivalent forms of the BktB protein of the present invention also include polypeptides and proteins wherein regions required for BktB-like catalytic activity are essentially
25 conserved or retained, but wherein non-critical regions may be modified by amino acid substitutions, deletions, or additions.

Due to the degeneracy of the genetic code, i.e., the existence of more than one codon for most of the amino acids naturally occurring in proteins,
30 genetically degenerate DNA (and RNA) sequences that contain the same essential genetic information as the DNA of SEQ ID NO:9 of the present

invention, and which encode the same amino acid sequence as that of *A. eutrophus* BktB, are encompassed by the present invention. Genetically degenerate forms of any of the other nucleic acid sequences discussed herein are encompassed by the present invention as well.

5

The nucleotide sequences described above are considered to possess a biological function substantially equivalent to that of the *A. eutrophus* BktB gene of the present invention if they encode peptides, polypeptides, or proteins having β -ketothiolase enzymatic activity differing from that of *A. eutrophus* BktB by about $\pm 30\%$ or less, preferably by about $\pm 20\%$ or less, and more preferably by about $\pm 10\%$ or less when assayed *in vivo* by complementation or by the enzymatic assays discussed above.

Based on amino acid sequence homology or sequence identity, one might attempt to predict the function of a protein of interest; however, this may not accurately reflect the activity of the protein, and it would therefore be necessary to assay the β -ketothiolase enzymatic activity thereof.

Other β -ketothiolases useful in the present invention may be derived from enzymes involved in lipid biosynthesis and β -oxidation which are not directly involved in PHA production.

Example 9

Production of PHBV in *E. coli* expressing *ilvA* and *bktB*

25

Genes required for the pathway diagrammed in figure 3 were assembled in recombinant *E. coli* in order to test the feasibility of producing PHBV from threonine in a pathway utilizing *ilvA* and *bktB* (both discussed above). *E. coli* DH12s (obtained from Gibco/BRL) was transformed with pMON25779 (harboring *A. eutrophus phbB* and *phbC* under LacI control) and pMON25822,

30

which expresses *bktB* from the *A. eutrophus bktB* promoter. The resulting recombinant *E. coli* strain expresses BktB β -ketothiolase, PhbB β -ketoacyl-CoA reductase, and PhbC PHA synthase.

pMON25779 was constructed as follows. pJM9131 (Kidwell et al., 1995) was digested with Stu I and the vector fragment was re-closed, thereby deleting a fragment of approximately 1 Kb that is required for production of the PhbA β -ketothiolase. The resulting plasmid, designated pMON25748, was digested with Bsa AI and Eco RI, and the fragment encoding a portion of *phbC* and the entire *phbB* gene was used to replace the Bsa AI-Eco RI fragment encoding a portion of *phbC* in pMON25629 (Figure 21), producing pMON25779. pMON25779 encodes *phbCB* under Ptac promoter control.

pMON25822 (Figure 25) contains a fragment of approximately 1.8 Kb encoding *A. eutrophus bktB* cloned into in the vector pBBR1MCS-3 (Kovach et al., 1994). pMON25822 was constructed as follows. pMON25765 is a pBluescript KS+ derivative that contains A 1.8 Kb Fragment encoding *bktB*. This plasmid is constructed by digesting pMON25728 (discussed above) partially with Sal I, then completely with Xho I, and cloning the resulting 1.8 Kb fragment into pBluescript KS+ that has been digested with Sal I and Xho I. The 1.8 Kb *bktB* fragment contains a total of 3 Sal I sites. pMON25765 was digested with Xho I and XbaI, and the 1.8 Kb fragment encoding *bktB* was ligated to pBBR1 MCS-3 digested with Xho I and XbaI.

To the strain harboring pMON25779 and pMON25822 was added either pSE280 (an expression vector lacking an *ilvA* gene: Brosius, 1989) or pMON25683 (expressing *ilvA466* from the LacI-regulated promoter of pSE280: figure 11). Each strain was grown to late log phase in 50 ml LB containing appropriate antibiotics, then the cells were pelleted by centrifugation, washed 2 times with M9 minimal salts, and resuspended in 100 ml M9 minimal media containing 0.5% glucose as the carbon source. The resuspended cells were split into two, 50 ml cultures, and threonine was added to one flask from each pair to a final concentration of 25 mM. The flasks were incubated overnight at 37°C with vigorous shaking. After about 18 hours, the

cells were concentrated by centrifugation, and polymer composition was analyzed as described above. The results are shown in Table 12.

Table 12
Production of PHBV from glucose in *E. coli*
expressing *ilvA* and *bktB*

| plasmid <i>ilvA</i> allele ¹ | Threonine added? | mol% hydroxyvalerate |
|---|------------------|----------------------|
| none | Y | 3.5 |
| none | N | N.D. ² |
| <i>ilvA466</i> ³ | Y | 31.4 |
| <i>ilvA466</i> ³ | N | 7.9 |

¹All strains are chromosomally *ilvA*⁺.

²N.D. = none detected.

³*ilvA466* contains the L481F mutation.

As demonstrated in Table 12, *E. coli* harboring *ilvA466* and *bktB* can make PHBV, even in the absence of added threonine. *E. coli* lacking a plasmid expressing *ilvA466* also produced some PHBV if threonine was added to the media. In this case, conversion of threonine to α -ketobutyrate presumably proceeds via the chromosomally-encoded IlvA protein.

These data demonstrate that substitution of *bktB* for *phbA* overcomes the metabolic block to PHBV formation in *E. coli* expressing only the *phb* operon (Table 3). These data also demonstrate that the PHBV copolymer can be produced *in vivo* from endogenous substrate pools if the cell expresses a partially-deregulated threonine deaminase and PHA biosynthesis enzymes having appropriate substrate flexibility.

Example 10

Production of P(3HB) in Seeds of Canola and Soybean

Poirier et al. (1992) and Nawrath et al. (1994) demonstrated the production of poly(3-hydroxybutyrate) (P(3HB)) homopolymer in leaves of transgenic *Arabidopsis* plants. Although this was the first demonstration of a plant-produced PHA, *Arabidopsis* is not an attractive commercial candidate for the large scale production of PHAs as this plant is primarily used as a research tool, and has no agronomic value. However, crops such as canola and soybean, for example, are excellent candidates for commercial PHA production in view of the vast acreage upon which they are grown. In addition, these crops are used for vegetable oil production, and significant pools of acetyl-CoA are present to support the oil biosynthesis that occurs in the seeds. As noted earlier, plant-produced PHAs have the potential to lower the current high cost of the microbially-produced PHAs by eliminating the fermentation costs and the expensive feedstocks required by microorganisms. For example, with properly timed and targeted expression of the PHB biosynthetic enzymes in seed plastids, one would expect to produce high levels (10%-20% fresh seed weight) of PHB polymer. Described below are the first definitive experimental results demonstrating PHA production in commercially useful plants, such as canola and soybean.

Transformation Vectors

Two *Agrobacterium* plant transformation vectors were constructed to introduce the entire PHB biosynthetic pathway from *A. eutrophus* into canola and soybean. The first plasmid, pMON25626 (Figure 23), comprises the genetic information for the expression of three genes in the transformed plant: *phbC*, *phbB* and CP4 EPSPS. Specifically, pMON25626 contains, beginning at the right border (RB) and continuing clockwise: the 7S β -conglycinin seed-specific promoter (Doyle et al., 1986; Slighton and Beachy, 1987) followed by the *Arabidopsis* small subunit of RUBP carboxylase chloroplast transit peptide

(Arab-SSU1A; Stark et al., 1992) translationally fused via an Nco I site to the *phbC* gene, followed by the polyadenylation and transcription termination signal contained in E 9 3' (Coruzzi, et al., 1984; Morelli et al., 1985). The vector continues with the 7S promoter, Arab-SSI1A, *phbB*, and E9 3' termination
5 signal, constructed as above. Additionally, pMON25626 contains the FMV promoter (Richine et al., 1987), followed by the petunia EPSPS (5-enol-pyruvylshikimate-3-phosphate synthase) chloroplast transit peptide (PEPSP; Shah et al., 1986) translationally fused to the CP4 EPSPS gene (PCT International Publication WO 92/04449; Padgett et al., 1996), and the *nos* 3'
10 polyadenylation and transcription termination signal (Fraley et al., 1983). The CP4 EPSPS gene was used as the selectable herbicide resistance marker ("glyphosate selection") in the transformation procedure (Shah et al., 1986). The CP4 EPSPS enzyme is a glyphosate-resistant form of the EPSPS enzyme involved in aromatic amino biosynthesis in plants and bacteria which catalyzes
15 the reversible reaction of shikimate-3-phosphate and phosphoenolpyruvate to produce 5-enolpyruvylshikimate-3-phosphate (Padgett et al., 1996).

The second plant transformation vector, pMON25638 (Figure 24), containing the *phbA* and the CP4 EPSPS genes, was constructed in a
mannersimilar to that of pMON25626. The promoters, chloroplast transit
20 peptides, and polyadenylation and transcription termination signals were identical to those described above for pMON25626. Both pMON25626 and pMON25638 contain the identical border sequence and replication origins and functions as described for pMON25668 of Example 5 in the section entitled
"Expression and Activity in Transformed Soybean Callus".

25

Transformation Protocols

pMON25626 and pMON25638 were introduced into *Agrobacterium tumefaciens* ABI (Koncz and Schell, 1986) by triparental mating (Ditta et al.,
30 1980), which was then used for the transformation of canola (Fry et al., 1987;

Radke et al., 1988) and soybean (Hinchee et al., 1988), with the modifications indicated below.

Canola Transformation

5

Plant Material

Stock plants were produced from seeds of the Westar variety planted in Metro Mix 350 and germinated in a growth chamber under a day temperature of 15°C, a night temperature of 10°C, a 16 hour day/8 hour night illumination period, a light intensity of 600 $\mu\text{En m}^{-2}\text{s}^{-1}$, and 50% relative humidity. Seedlings were subirrigated with water daily, and soaked with a 15-30-15 nutrient solution every other day for one hour. At three weeks, seedlings were transferred into 6" pots. Five week old plants were harvested once the plants bolted, but prior to flowering (plants with up to three flowers can be employed, however). The leaves and buds were removed from the stem, and the 4-5 inches of stem just below the flower buds were used as the explant tissue source. Just prior to inoculation, the stems were sterilized by soaking in 70% ethanol for 1 min, 38% Chlorox (4% sodium hypochlorite) for 20 min, rinsing two times in sterile deionized water, and soaking in two tablespoons of Captan (Captan 50-WP, ICI Ag Products) plus 500 mls sterile water for 15 min.

Preparation of *Agrobacterium*

Five to 7 days prior to inoculation, *Agrobacterium* was streaked from a frozen glycerol stock onto an LB plate (1.5% agar) containing 100 mg/l spectinomycin, 100 mg/l streptomycin, 25 mg/l chloramphenicol, and 50 mg/l kanamycin (denoted LBSSCK). Two days before inoculation day, a 10 μl loop of *Agrobacterium* was placed into a tube containing 2 mls of LBSSCK and placed on a rotator overnight at 22-28°C. The day before inoculation, the *Agrobacterium*

was subcultured by placing 200 μ l in a tube containing 2 ml of fresh LBSSCK, which was placed on a rotator overnight. On the day of inoculation, the *Agrobacterium* was diluted 1:10 with MS liquid medium (Murashige and Skoog, 1962) to an OD₆₆₀ of 0.2-0.4.

5

Explant Inoculation

Sterilized stems were cut into 0.6 cm segments (0.3-1.5 cm segments can be used), noting their basal orientation. Explants were inoculated for five minutes in a square Petri plate (100X15 mm) with the 1:10 dilution of *Agrobacterium*. Five mls of *Agrobacterium* solution were added to five stems by pipetting the *Agrobacterium* directly on top of the explants. After five minutes, the *Agrobacterium* solution was aspirated off the explants. The stem explants were then cultured in the basal-side down orientation for an optimal shoot regeneration response on the co-culture plates. Co-culture plates (100X15 mm) contained 1/10 MS salts (this can range from about 1/10 to full strength; Gibco, 500-1117EH), 1X B5 vitamins (Sigma, G-2519), 0.5 mg/l 6-benzylamino-purine (this can range from about 0.1-2 mg/l), 3% sucrose (this can range from about 1-6%), pH 5.7, solidified with 0.9% agar, covered with 2 ml TXD liquid medium (Horsch et al., 1985) onto which an 8.5 cm piece of sterile Whatman qualitative grade filter paper was placed. Excess *Agrobacterium* present on the stem explants placed on the filter paper was blotted off using another piece of sterile 8.5 cm filter paper. The co-culture plates were placed in clear plastic bags which were slit on the sides to permit air exchange, and which were incubated in a warm room at 25°C under 24 hours continuous cool white light (40 μ En m⁻²s⁻¹).

Tissue Selection and Regeneration

After two days, the stem explants were moved onto MS medium containing 500 mg/l ticarcillin, 50 mg/l cefotaxime, and 1 mg/l 6-benzylaminopurine for a three day delay period. Plates were again placed in slit, clear plastic bags which were placed in the warm room. After a three day delay

5 period, stem explants were moved onto glyphosate selection medium containing MS salts, B5 vitamins, 0.1 mM glyphosate (this can range from about 0.025-0.2 mM), 500 mg/l ticarcillin (this can range from about 250-750 mg/l), 50 mg/l cefotaxime (this can range from about 25-100 mg/l), and 1 mg/l 6-benzylaminopurine (this can range from about 0.1-4 mg/l) for three weeks. After three

10 weeks, the stem explants were moved onto glyphosate selection medium containing MS salts, B5 vitamins, 0.1 mM glyphosate (this can range from about 0.025-0.2 mM), 500 mg/l ticarcillin (this can range from about 250-750 mg/l), 50 mg/l cefotaxime (this can range from about 25-100 mg/l), and 1 mg/l 6-benzylaminopurine (this can range from about 0.1-4 mg/l), plus 0.5 mg/l

15 gibberellic acid A3 (this can range from about 0.1-2 mg/l), which enhances shoot elongation, for another three week period. After these six weeks on glyphosate selection medium, normally developing green shoots were excised from the stem explants. Shoots (4-5 per plate) were placed in rooting medium (1/10-full strength MS salts, Staba vitamins (Staba, 1969), 3% sucrose (this can range

20 from about 1-6%), 500 mg/l ticarcillin (this can range from about 250-750 mg/l), 50 mg/l cefotaxime (this can range from about 25-100 mg/l), and 2 mg/l indolebutyric acid (this can range from about 0.5-3 mg/l), pH 5.7, solidified with 0.9% agar. Root development began to occur as early as one week after shoots were placed on rooting medium. At the two week timepoint, shoots having a

25 large root base were moved into 2 1/2" pots containing Metro Mix 350 (Hummert Co., St. Louis, MO). Flats were covered with clear plastic domes (Hummert Co., St. Louis) so the shoots could elongate. Flats containing R₀ plants were placed in a growth chamber under the same conditions as described above for stock plant growth. After 3-4 days, the domes were cracked in order

30 to harden off the plants under the following conditions: Temperature: 20°C day/15°C night; Photoperiod: 16 hr light/8 hr dark; Light intensity: 450 µEn

m-2s-1; Relative humidity: 70%; Fertilizer: 15-16-17 Peter's Solution (200 ppm nitrogen). Hardened plants were grown for approximately 14 weeks under the same conditions, at which time seeds were collected. Cross-pollination was prevented by bagging the plants at bolting time.

5

This protocol results in transformation efficiencies (defined as the number of confirmed transgenics/ the number of explants inoculated, expressed as a percentage) as high as 35-40%. This is a significant improvement over the protocol using kanamycin selection (Fry et al., 1987).

10

Soybean Transformation

Plant Material

15 Seeds of soybean variety A3237 were surface sterilized by rinsing them in dilute Tween 20 (polyoxyethylenesorbitan monolaurate) for 30 seconds, followed by rinsing under running tap water for approximately two min. The seeds were then rinsed in 80% ethanol, and then agitated in freshly made 50% Chlorox (5.25% sodium hypochlorite) containing Tween 20 for 15 min. The
20 seeds were then completely rinsed with five rinses of sterile distilled water. They were then placed in a saturated Captan and/or Benylate slurry for 2-30 min to control fungus infestation.

25 Sterilized seeds were then placed on 0.7% purified agar-solidified B5 basal medium (Gamborg et al., 1968) for germination (approximately 15 seeds per plate). The Petri dishes were placed in a plastic bag slit on the sides to permit air exchange, and incubated in a culture room under 18-20 hrs light ($60 \mu\text{En m}^{-2}\text{s}^{-1}$), 4-6 hrs dark, at 25°C , for 5-6 days. After this incubation, the germinated seeds were placed in a cold room or refrigerator ($0-10^{\circ}\text{C}$; average
30 temperature of 4°C) for at least 24 hrs prior to explanting.

Preparation of *Agrobacterium*

Agrobacterium strains to be used for transformation were prepared as follows. Bacteria were streaked from frozen glycerol stocks onto LBSCK plates containing 1.5% agar-solidified LB medium plus 100 mg/l of spectinomycin, 25 mg/l of chloramphenicol, and 50 mg/l of kanamycin. The bacteria can be incubated at room temperature or in an incubator at 27°C for 2-4 days. Prior to preparing the *Agrobacterium* inoculum, a fresh plate of *Agrobacterium* was streaked from the first plate 2-3 days prior to growth on liquid medium. One to two days prior to the inoculation of soybean explants, one loop of bacteria was transferred from a freshly streaked plate into a culture tube containing 2 ml of YEP medium containing 10 g/l peptone, 10 g/l yeast extract, 5 g/l NaCl, 100 mg/l spectinomycin, 25 mg/l chloramphenicol, and 50 mg/l kanamycin. Larger volumes of bacteria can be grown using the same basic formula of one loop of bacteria per 2 ml of YEP. The tube containing the bacteria in YEP was vortexed to disperse the clump of bacteria, and placed on a rotator. For a one day culture, the bacteria can be started at about 7:00 a.m.; for a two day culture, the bacteria can be started later in the day and allowed to grow overnight. The afternoon prior to inoculating the explants, 4-6 mls (2-3 tubes) of the bacterial culture were added to 50 mls of AB minimal salts medium (Chilton et al., 1974) containing the same concentrations of spectinomycin, chloramphenicol, and kanamycin as in the LBSCK medium, in sterile 250 ml flasks. This culture was grown on a shaker overnight at 28°C. The bacteria were pelleted by centrifugation and the pellet was resuspended to an OD₆₆₀ of 0.25-1.0 with the following medium: 1/10 B5 salts (this can range from about 1/10 to full strength), 1/10 B5 vitamins (this can range from about 1/10 to full strength), 3% sucrose or glucose (this can range from about 0.5-6% sucrose or glucose), 7.5µM 6-benzyl-aminopurine (this can range from about 2.5-20 µM), 200 µM acetosyringone (this can range from about 50-300 µM), 1mM galacturonic acid (this can range from about 0.1-2mM), 0.25 mg/l gibberellic

acid (GA3) (this can range from 0-0.5 mg/l), and 20mM MES, pH 5.4 (the pH can range from about 5.2-6.0).

Explant Inoculation

5

Explants were prepared by removing the seed coat from the germinated seedlings and cutting the hypocotyl at approximately 0.5 cm or more from the cotyledons (one cm is preferred). The lower portion of the hypocotyl and root axis was discarded. The cotyledons and remaining hypocotyl were completely split by making an incision down the middle of the hypocotyl and then bending the halves apart so that they separated from one another. The primary leaves and primary shoot meristem were removed. The region of the cotyledon near the axillary bud was wounded multiple times (anywhere from 3-15 times) using a scalpel blade, the score marks being placed longitudinally with respect to the embryo axis. The axillary bud can be damaged in the process, but this is not required. Approximately 40-80 explants were prepared and added to a single, dry Petri dish. Approximately 10 mls of the bacterial inoculum were added to just cover the explants. The explants remained in contact with the *Agrobacterium* solution for 30 min. The *Agrobacterium* solution was then removed from the explants which were briefly blotted on sterile Whatman filter paper prior to being placed flat (adaxial) side down onto co-culture plates. Co-culture plates were prepared by adding 4-5 mls of the bacterial dilution medium additionally containing 3% sucrose, 1mM galacturonic acid, and 200µM acetosyringone to 1-2 layers of sterile Whatman filter paper in a 100X15mm Petri dish. The co-culture medium can contain a mixture of 0.5-6% glucose or 0.5 -6% sucrose (1-3% of either being preferred), with or without 0.1-10mM galacturonic acid (1mM being preferred), with or without 50-300µM acetosyringone (100-200µM being preferred). The co-culture medium was solidified with 0.8% washed agar (Sigma, A 8678).

30

Tissue Selection and Regeneration

The explants were co-cultured with the *Agrobacterium* in a culture room at 20-23°C under an 18-20 hr light/4-6 hr dark photoperiod (co-culturing can be carried out from about 18-26°C). Co-culture lasted for 2-4 days. After co-culture, the explants were washed in wash medium containing 1/10 B5 salts (this can range from about 1/10 to full strength), 1/10 B5 vitamins (this can range from about 1/10 to full strength), 7.5µM 6-benzylaminopurine (this can range from about 2.5-20µM), pH 5.6 (the pH can range from about 5.2-6.0), 500 mg/l ticarcillin (this can range from about 250-750 mg/l), and 100 mg/l cefotaxime (this can range from about 25-200 mg/l) .

The washed explants were cultured on a culture medium containing B5-basal salts and vitamins, 7.5µM 6-benzylaminopurine (this can range from about 2.5 µM-20 µM), 500 mg/l ticarcillin (this can range from about 250-750 mg/l), 100 mg/l cefotaxime (this can range from about 25-200 mg/l), and 0.075-0.1 mM glyphosate (this can range from about 0.025-0.4 mM). The plates were sealed with white 3M porous tape and placed in a culture room or incubator at 24-26°C under an 18-20 hr light/4-6 hr dark cycle at 20-80 µEn m⁻²s⁻¹. Subsequent subcultures were made every 2-3 weeks.

At two to four weeks, the cultures were transferred to MSB5 medium (Sigma, M 0404 or Gibco, 500-117EH plus Sigma, G2519) or B5 basal medium plus 1 mg/l zeatin riboside (this can range from about 0-5 mg/l), 0.5 mg/l gibberillic acid (GA3) (this can range from about 0-2 mg/l) , 0.1 mg/l indoleacetic acid (this can range from about 0-1 mg/l), 2.5µM 6-benzylaminopurine (this can range from about 0-5 µM), 500 mg/l ticarcillin (this can range from about 250-750 mg/l), 100 mg/l cefotaxime (this can range from about 25-200 mg/l), and 0.075mM glyphosate (this can range from about 0.025-0.2 mM). Additional B5 micronutrients (up to four times the standard concentration of each

micronutrient alone or in various combinations with the others) and 2 gm/l proline (this can range from about 0-2 gm/l) can be added to this medium.

- At the four to six week time point, the petiole/hypocotyl tissue and
- 5 cotyledons, as well as any dead or dying material, i.e., any non-regenerating tissues, were removed (such material can generally be removed between 4-9 weeks). The regenerating cultures were transferred to 0.8% washed agar-solidified elongation medium comprising MSB5 medium or B5 basal medium plus 1 mg/l zeatin riboside (this can range from about 0-5 mg/l), 0.5 mg/l
 - 10 gibberillic acid (this can range from about 0-2 mg/l), 0.1 mg/l indoleacetic acid (this can range from about 0-1 mg/l), 500 mg/l ticarcillin (this can range from about 250-750 mg/l), 100 mg/l cefotaxime (this can range from about 25-200 mg/l), and 0.05mM glyphosate (this can range from about 0.025-0.2 mM), and again placed in a culture room or incubator at 24-26°C under an 18-20 hr
 - 15 light/4-6 hr dark cycle at 20-80 $\mu\text{En m}^{-2}\text{s}^{-1}$. Elongation medium can contain about 0.25-2 mg/l zeatin riboside, 0.01-1 mg/l indoleacetic acid, and 0.1-5 mg/l gibberellic acid (GA3). Cultures were transferred every three weeks to the same medium. Identification of putative A3237 transgenics (elongating, normal appearing shoots) required approximately 8-20 weeks.
 - 20 Shoots were rooted on 0.7% purified agar-solidified one-half or full strength MSB5 medium or one-half or full strength B5 basal medium containing 500 mg/l ticarcillin (this can range from about 0-500 mg/l), 100 mg/l cefotaxime (this can range from about 0-100 mg/l), and 1 mg/l indolebutyric acid (this can range from about 0.1-2 mg/l) or naphthaleneacetic acid (this can
 - 25 range from about 0.05-2 mg/l), with 0-50 mg/l glutamine and 0-50 mg/l asparagine at 24-26°C under an 18-20 hr light/4-6 hr dark cycle for 2-6 weeks. Rooted shoots were placed in 2" pots containing moistened MetroMix 350, and kept enclosed in magenta boxes until acclimatized at 24-26°C under an 18-20 hr light/4-6 hr dark cycle (20-80 $\mu\text{En m}^{-2}\text{s}^{-1}$). Shoots were hardened off for 3-4
 - 30 days after cracking the lids under the following conditions: Photoperiod: 18-20 hrs light/4-6 hrs dark; Light intensity: 20-80 $\mu\text{En m}^{-2}\text{s}^{-1}$; Temperature: 24-

26°C. Hardened plants were grown for approximately 3 weeks under the following conditions: Photoperiod: 12 hr light/12 hr dark; Light intensity: 450 $\mu\text{En m}^{-2}\text{s}^{-1}$; Relative humidity: 70%; Temperature: 26°C day/21°C night. Transformation was confirmed by detection of expression of the CP4 gene by
 5 an ELISA assay. CP4-positive plants were subsequently grown under the following conditions: Photoperiod: 12 hr light/12 hr dark; Light intensity: 450 $\mu\text{En m}^{-2}\text{s}^{-1}$; Relative humidity: 70%; Temperature: 26°C day/21°C night; Fertilizer: 15-16-17 Peter's Solution (200 ppm nitrogen). Plants were grown for approximately 11 weeks, at which time seed was collected.

10

Glyphosate Selection

Glyphosate (0.05 mM-0.1 mM) was employed as a selectable marker (Hinchee et al., 1994) for both canola and soybean. Leaves of glyphosate-
 15 resistant canola and soybean transformants (designated R_0 generation) were screened for CP4 EPSPS expression by ELISA (Padgett et al., 1995). Seeds from R_0 CP4-positive plants were assayed enzymatically for either the PhbA β -ketothiolase or PhbB reductase, depending upon the plasmid used for transformation (data not shown). PHA synthase was not measured
 20 enzymatically; however, this gene is carried on the same genetic insert as the *phbB* reductase (see Figure 23), and plants containing PhbB activity are likely to contain PhbC activity as well. Soybean and canola seed extracts were prepared by grinding the seeds to a fine powder in liquid N_2 , washing with acetone three times, and filtering through Whatman 3MM paper. The seed
 25 extracts were dried at 37°C, one ml of extraction buffer (100mM KPi, pH7.4, 5mM MgCl_2 , 1mM EGTA, 5mM DTT, 10% glycerol) was added, the mixture was vortexed, and the insoluble fraction was pelleted. The extracts were used for enzyme assays and to detect all PHB biosynthetic enzymes by Western blot analysis according to Nawrath et al. (1994). Immunoreactive bands
 30 corresponding to each of these enzymes were detected in extracts of seeds of both plants (data not shown).

Since the CP4 EPSPS gene is genetically linked to the β -ketothiolase gene in pMON25638, and the reductase and PHA synthase genes in pMON25626, glyphosate spray selection (Padgett et al., 1995) was employed to identify plants retaining the CP4 EPSPS locus and respective linked PHB biosynthetic genes. Plants surviving the glyphosate spray contain the CP4 EPSPS gene and the linked PHB biosynthetic genes.

Several positive reductase- (pMON25626) and β -ketothiolase- (pMON25638) expressing plants were tested for segregation by spraying glyphosate on approximately 40 siblings of the R_1 generation (derived by self-pollination of R_0 plants). Any plant that lost the CP4 EPSPS gene due to genetic segregation will die when sprayed with glyphosate. Plant lines that segregated three CP4 positive:one CP4 negative (3:1, indicating the presence of only a single genetic locus) were retained for production of homozygous plants for the respective β -ketothiolase and reductase loci. For each independent line that segregated 3:1, approximately ten R_1 sibling plants per line were retained for seed production. Seeds from each of the siblings (designated R_2 generation) were planted, and approximately 40 plants per line were sprayed with glyphosate. Lines in which all R_2 siblings survived the glyphosate spray were considered to be homozygous for the respective loci.

In order to assemble the entire PHB biosynthetic pathway in a single plant, homozygous plants derived from transformed canola or soybean containing pMON25626 (PhbC and PhbB) were crossed with homozygous plants containing pMON25638 (PhbA). Seeds from the crosses (designated F_1) were assayed for the presence of the β -ketothiolase and reductase to confirm successful crossing (data not shown). At this point in the breeding cycle, the plants are not homozygous for all three genes. The locus containing the β -

ketothiolase gene segregates independently from the locus containing the reductase and synthase genes. The segregation ratio in the F_1 generation would follow a typical Mendelian pattern of inheritance. Five F_1 progeny seeds were planted, and seeds obtained therefrom (designated F_2) were collected and
5 assayed for the presence of PHB as described below.

Isolation and Characterization of Poly(3-Hydroxybutyrate) from Transgenic Canola Seeds

10 40 g of F_2 transgenic canola seeds were crushed in a mechanical benchtop crusher (Cemotec 1090 Sample Mill, Grinding Disc set to position 1) and then transferred to cellulose extraction thimbles (Whatman, Single thickness, 26mm external diameter x 60mm external length). Extraction was carried out in these thimbles under refluxing chloroform at 80°C using a Soxtec
15 Extraction System (Tecator).

After approximately 12 hours of extraction, the extract was collected in a flask, and the chloroform evaporated in a fume-hood under a stream of air/nitrogen, leaving behind canola oil and an insoluble, gel-like solid residue. To
20 this mixture, 50 ml of hexanes were added, and the mixture was stirred for 5 minutes to dissolve the oil. The mixture was then centrifuged for 5 minutes at 1,700 x g, producing a white pellet. The supernatant was decanted off, and the pellet redissolved in 10 ml chloroform. Upon stirring this solution with 100 ml hexanes, a white precipitate was obtained. This precipitate was recovered by
25 centrifugation, washed twice with 30 ml hexanes to remove the residual oil, and dried under a stream of nitrogen for two hours at room temperature, and under vacuum for three hours at approximately 40-50°C.

Structural analysis of a 3 mg/ml solution of the precipitate in deuterated-
30 chloroform (99.8% D, Aldrich Chemical Co., USA) by a 300-MHz ^1H -NMR

performed on a Varian VX-300 spectrometer confirmed it to be pure Poly(3-hydroxybutyrate) (P(3HB)) (data not shown).

Molecular weight data were obtained at 35°C using a Waters Millennium
5 GPC System and a 410 Differential Refractometer with Waters Ultrastyrigel
Columns of pore sizes 105 Å, 104 Å, 103 Å and 500 Å connected in series.
Chloroform was used as eluent at a flow rate of 1 ml/min, and a sample
concentration of 6 mg/ml was used. Polystyrene standards with a low
polydispersity (Aldrich Chemical Company, USA) were used to prepare a
10 calibration curve. The results are shown in Table 13.

15

20

TABLE 13
Production of P(3HB) in Transgenic Canola Seeds

| Weight of canola seeds | Weight of PHB obtained | % PHB (with respect to seed weight) | Mw ¹ (Daltons) | Mn ² (Daltons) | Polydispersity (Mw/Mn) |
|------------------------------|------------------------------|--|------------------------------|------------------------------|---------------------------|
| 40 g | 0.2 g | 0.5% | 686,300 | 276,800 | 2.5 |

¹Mw: Weight average molecular weight.

25 ²Mn: Number average molecular weight.

Isolation and Characterization of Poly(3-Hydroxybutyrate) from Transgenic Soybean Seeds

- 5 The extraction procedure described above for canola seeds was also employed for transgenic soybean seeds, using 10 g of F₂ generation crushed seeds. The results are as summarized in Table 14.

10

TABLE 14
Production of P(3HB) in Transgenic Soybean Seeds

| Weight of soybean seeds | Weight of PHB obtained | % PHB (with respect to seed weight) | Mw ¹ (Daltons) | Mn ² (Daltons) | Polydispersity (Mw/Mn) |
|-------------------------|------------------------|-------------------------------------|---------------------------|---------------------------|------------------------|
| 10 g | 0.0015 g | 0.015% | 209,700 | 99,400 | 2.1 |

¹Mw: Weight average molecular weight.

²Mn: Number average molecular weight.

- 15 The data presented in Tables 13 and 14, respectively, confirm the biosynthesis of P(3HB) in canola seeds at a level of at least 0.5% of total seed weight, as well as the biosynthesis of P(3HB) in soybean seeds at a level of at least 0.015% of the total seed weight, by transformation and breeding employing plastid-targeted, seed-specific expression of the *A. eutrophus* PHB
- 20 biosynthetic enzymes. These results represent the first definitive demonstration of the production of PHA in oilseeds of transgenic commercial crops.

Example 11

25

Optimization of P(3HB-co-3HV) Copolymer Production in Plant and Bacterial Cells

The ability to biosynthesize PHAs such as P(3HB-co-3HV) copolymer can be conferred upon heterologous host cells by introduction therein of appropriate enzyme-encoding nucleic acids in such a manner that they express enzymatically active products. Introduction into bacterial and plant cells of genetic constructs containing chromosomal DNAs, plasmid DNAs, cDNAs, and synthetic DNAs encoding PHA biosynthetic enzymes, in various combinations with the enzymes discussed above, wherein such nucleic acids are operably linked to appropriate regulatory regions for expression depending upon the host cell, enables such cells to biosynthesize and accumulate PHAs. In bacteria for example, Slater et al. (1988) and Schubert et al. (1988) have demonstrated that *E. coli* transformed with and expressing the *A. eutrophus phbA*, *phbB*, and *phbC* genes produces significant levels of PHB. In plants, Poirier et al. (1992) have shown that *Arabidopsis* expressing these genes produces the PHB homopolymer. Higher levels of PHB were obtained by targeting the Phb enzymes to the leaf plastids (Nawrath et al., 1994).

In view of the ability of heterologous bacterial and plant host cells to produce PHB polyhydroxyalkanoate upon introduction and expression therein of DNAs encoding the appropriate PHB biosynthetic enzymes, it is similarly expected that P(3HB-co-3HV) copolymer can be produced in bacteria and plants by expressing therein appropriate combinations of PHA-biosynthetic and other enzymes as discussed above. As the levels of acetyl-CoA required for copolymer biosynthesis in bacteria and plant cells appear to be non-limiting (Nawrath et. al., 1994), no further manipulation of cellular metabolism with respect to this precursor is likely to be necessary. Insuring the presence of sufficient pools of propionyl-CoA required for C4/C5 copolymer production may require introduction into, and overexpression in, host cells of various combinations of wild-type or deregulated aspartate kinase, homoserine dehydrogenase, threonine synthase, wild-type or deregulated threonine

deaminase, α -ketoacid dehydrogenase E1, E2, and E3 subunits, pyruvate oxidase, and acyl-CoA synthetase enzymes.

Thus, DNA encoding the following enzymes can be introduced into and
5 expressed in plants and bacteria in which P(3HB-co-3HV) copolymer is desired:
an appropriate β -ketothiolase or combination of β -ketothiolases, a β -ketoacyl-
CoA reductase, and a PHA synthase. If necessary, DNA encoding a wild-type
or deregulated threonine deaminase can also be introduced into and expressed in
these organisms. The β -ketothiolase can be the BktB β -ketothiolase of *A.*
10 *eutrophus*, or one similar thereto, that can condense two molecules of acetyl-
CoA to produce acetoacetyl-CoA, and that can also condense acetyl-CoA and
propionyl-CoA to produce β -ketovaleryl-CoA. Alternatively, a combination of
 β -ketothiolases can be employed, i.e., a β -ketothiolase capable of condensing
two molecules of acetyl-CoA to produce acetoacetyl-CoA, and a β -ketothiolase
15 capable of condensing acetyl-CoA and propionyl-CoA to produce β -ketovaleryl-
CoA. The β -ketoacyl-CoA reductase can be one such as that encoded by the
A. eutrophus phbB gene; the PHA synthase can be one such as that from *A.*
eutrophus as well. *A. eutrophus* is only one of many microorganisms that
produce PHAs, and DNA (genomic, plasmid, cDNA, or synthetic DNA) encoding
20 any of the foregoing enzymes can be derived from any other PHA-producing
organism containing enzymes having the same or similar enzymatic activity as
the *A. eutrophus* enzymes, as discussed above in Example 8 in the section
entitled "Other β -Ketothiolases, β -Ketoacyl-CoA Reductases, and PHA
Synthases for the Production of P(3HB-co-3HV) Copolymer."

25

As described in Examples 1-8, there are a number of different methods
by which the levels of α -ketobutyrate, propionyl-CoA, and ultimately 3-

hydroxy-valeryl-CoA which can be utilized in PHA synthesis can be increased in heterologous bacterial and plant host cells. These are summarized below.

Overexpression of Threonine Deaminase

5

The simplest method of increasing the level of α -ketobutyrate in a bacterial or plant host cell involves overexpressing a wild-type or deregulated threonine deaminase in such cells. In plants, this enzyme can be expressed cytoplasmically, targeted to the plastids using appropriate chloroplast transit peptides, or expressed directly within these organelles by plastid transformation. These strategies are discussed in more detail below.

Overexpression of Aspartate Kinase and Threonine Deaminase

15 Overexpression of a wild-type or deregulated aspartate kinase will result in an increase in intracellular levels of L-threonine. Deregulated aspartate kinases have been described, for example, in *E. coli* (Boy et al., 1979; Karchi et al., 1993), *C. lactofermentum* (Jetten et al., 1995), barley (Bright et al., 1982), maize (Dotson et al., 1990; Hibberd et al., 1980), carrot (Cattoir-Reynaerts et al., 1983), and tobacco (Frankard et al., 1991). Expression of these enzymes in the cytoplasm, targeting of these enzymes to the plastids of plant cells, or expression of these enzymes in plastids following plastid transformation, will result in an increase in L-threonine therein. Simultaneous overexpression of a wild-type or deregulated threonine deaminase in bacterial cells, or in the cytoplasm or plastids of plant cells, will result in conversion of this excess L-threonine to α -ketobutyrate, which will then lead to increased levels of propionyl-CoA and then 3-hydroxyvaleryl-CoA, which can be employed in P(3HB-co-3HV) copolymer biosynthesis.

Overexpression of Aspartate Kinase, Homoserine Dehydrogenase, and Threonine Deaminase

An overexpressed homoserine dehydrogenase can be used in conjunction with a wild-type or deregulated aspartate kinase to increase further the level of threonine in bacterial cells, or in plant cells and plastids by employing

- 5 appropriate chloroplast transit peptides or chloroplast transformation. The excess threonine can then be directed to 3-hydroxyvaleryl-CoA synthesis via α -ketobutyrate by an overexpressed wild-type or deregulated threonine deaminase, which can be targeted to or expressed in plastids in plant cells.

10 **Overexpression of Threonine Deaminase, Pyruvate Oxidase, and Acyl-CoA Synthetase**

- α -ketobutyrate levels can be enhanced in bacterial cells and plant plastids (via chloroplast transit peptide targeting or plastid transformation) by
- 15 overexpressing a wild-type or deregulated threonine deaminase. The excess α -ketobutyrate can then be converted to propionate and CO₂ by an overexpressed pyruvate oxidase, for example, *E. coli* PoxB. Endogenous or overexpressed bacterial or plastid acyl-CoA synthetases will then activate the free propionate to produce propionyl-CoA, which can then be used by plants
 - 20 expressing other appropriate genes (for example, isoleucine-deregulated threonine deaminase and BktB β -ketothiolase) to form the 3-hydroxyvaleryl-CoA monomer employed in P(3HB-co-3HV) copolymer synthesis.

25 **Overexpression of Aspartate Kinase, Homoserine Dehydrogenase, Threonine Deaminase, Pyruvate Oxidase, and Acyl-CoA Synthetase**

- A wild-type or deregulated threonine deaminase, aspartate kinase, homoserine dehydrogenase, pyruvate oxidase, and an acyl-CoA synthetase, each having the effect variously described above, can be simultaneously
- 30 overexpressed in bacterial cells or plant cells and plastids (via the use of a

chloroplast transit peptide or plastid transformation) to provide increased levels of propionyl-CoA and, eventually, 3-hydroxyvaleryl-CoA, which can be incorporated into P(3HB-co-3HV) copolymer in the presence of β -ketothiolase, acetoacetyl-CoA reductase, and PHA synthase.

5

Overexpression of Aspartate Kinase, Homoserine Dehydrogenase, Threonine Synthase, Threonine Deaminase, Pyruvate Oxidase, and Acyl-CoA Synthetase

- 10 In addition to aspartate kinase, homoserine dehydrogenase, threonine deaminase, pyruvate oxidase, an acyl-CoA synthetase, and threonine synthase can also be introduced into plant and bacterial cells to provide increased levels of propionyl-CoA and, eventually, 3-hydroxyvaleryl-CoA, which can be incorporated into P(3HB-co-3HV) copolymer in the presence of β -ketothiolase,
- 15 acetoacetyl-CoA reductase, and PHA synthase.

20 **Overexpression of a Branched-Chain α -Ketoacid Dehydrogenase Complex E1 Subunit**

As discussed in Example 6, pyruvate dehydrogenase complex catalyzes the oxidative decarboxylation of α -ketobutyrate to form propionyl-CoA.

- 25 Pyruvate dehydrogenase complex is a multienzyme complex containing three activities: a pyruvate decarboxylase (E1); a dihydrolipoyl transacetylase (E2); and a dihydrolipoyl dehydrogenase (E3).

Overexpression of a branched-chain α -ketoacid dehydrogenase E1

30 subunit having improved binding and decarboxylating activity with

α -ketobutyrate as compared to the naturally occurring E1 subunit may effectively compete with the endogenous bacterial or plant E1 subunit. This overexpressed E1 subunit will combine with the endogenous pyruvate dehydrogenase E2E3 subcomplex to create a functional hybrid complex

- 5 capable of turning over α -ketobutyrate to propionyl-CoA, which can then be metabolized to 3-hydroxyvaleryl-CoA for incorporation into P(3HB-co-3HV) copolymer.

- Alternatively, both the E1 and E2 components of a branched-
10 chain α -ketoacid dehydrogenase complex that has significant activity with α -ketobutyrate can be overexpressed. Overexpression of the dihydrolipoyl dehydrogenase E3 subunit may not be necessary as this subunit is common to all α -ketoacid dehydrogenase complexes. However, if the endogenous E3 levels are not sufficient for E1E2 of endogenous PDC and the overexpressed E1E2 of
15 a branched-chain α -ketoacid dehydrogenase complex, then it too can be overexpressed.

- Overexpression of an α -ketoacid dehydrogenase complex E1 subunit having greater binding and decarboxylating activity with α -ketobutyrate than
20 the naturally occurring E1 subunit in bacterial cells or plant plastids, or overexpression of both E1 and E2 components as discussed above, can be employed in conjunction with any of the other methods discussed herein as a means for increasing the amounts of propionyl-CoA and subsequently 3-hydroxyvaleryl-CoA available for P(3HB-co-3HV) copolymer synthesis.
25

Production of Transformed Bacteria and Transgenic Plants
Capable of Producing P(3HB-co-3HV) Copolymer

PHA synthesis in bacteria and plants can be optimized in accordance with the present invention by expressing DNAs encoding β -ketothiolase, β -acyl-CoA reductase, and PHA synthase in conjunction with various combinations of precursor-producing enzymes, as discussed in the foregoing Examples.

5 Methods therefor are discussed below.

Bacterial Vectors

Methods for incorporating all the genes discussed herein into
10 transformation/expression vector constructs and introducing these constructs into bacterial and plant host cells to produce PHAs in such cells are well known in the art. Poirier et al. (1995) have recently provided an extensive review of progress in this area. In general, such vector constructs comprise assemblies of DNA fragments operatively linked in a functional manner such that they drive
15 the expression of the structural DNA sequences contained therein. These vector constructs usually contain a promoter that functions in the selected host cell, along with any other necessary regulatory regions such as ribosome binding sites, transcription terminators, 3' non-translated polyadenylation signals, etc., linked together in an operable manner, as well as selectable
20 markers (Sambrook et al., 1989; Ausubel et al. 1989). Vectors for bacterial cloning have been reviewed in Rodriguez et al. (1988).

Bacterial Transformation

25 Such vectors can be introduced into bacterial cells by calcium chloride/heat shock treatment or electroporation. Transformed host cells can subsequently be selected on selective media, cultured in an appropriate medium for a time and under conditions conducive to the production of PHA, and the PHA can then be recovered. Representative methods have been described by
30 Slater et al. (1988); Slater et al. (1992); Zhang et al. (1994); and Kidwell et al. (1995).

Bacterial and Other Host Cells

Useful host organisms for PHA polymer production include

- 5 Actinomycetes (e.g., *Streptomyces* sp. and *Nocardia* sp.); bacteria (e.g., *Alcaligenes* (e.g., *A. eutrophus*), *Bacillus cereus*, *B. subtilis*, *B. licheniformis*, *B. megaterium*, *Escherichia coli*, *Klebsiella* (e.g., *K. aerogenes* and *K. oxytoca*), *Lactobacillus*, *Methylobacter*, *Pseudomonas* (e.g., *P. putida* and *P. fluorescens*); fungi (e.g., *Aspergillus*, *Cephalosporium*, and *Penicillium*); and yeast (e.g.,
- 10 *Saccharomyces*, *Rhodotorula*, *Candida*, *Hansenula*, and *Pichia*).

- Organisms capable of overproducing threonine either naturally, via chemical or physical mutagenesis, or via recombinant DNA methodology (Jetten & Sinskey, 1995), can be used in the present invention for the
- 15 production of P(3HB-co-3HV) copolymer. DNAs selected from the group consisting of DNA encoding a wild-type or deregulated threonine deaminase, a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA, a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to form β -ketovaleryl-CoA, e.g., BktB, a β -ketoacyl-CoA
 - 20 reductase capable of converting a β -ketoacyl-CoA to its corresponding β -hydroxyacyl-CoA, e.g., PhbB, and a PHA synthase capable of producing P(3HB-co-3HV) copolymer from the β -hydroxyacyl-CoAs, e.g., PhbC, or combinations thereof, can be introduced into these organisms in order to impart P(3HB-co-3HV) copolymer production capability thereto.

25

Organisms that are capable of producing propionate or propionyl-CoA and other odd-chain substrates from simple precursors such as glucose, lactate, etc., are also useful for producing P(3HB-co-3HV) copolymer. For example, DNAs encoding a β -ketothiolase capable of condensing two molecules

of acetyl-CoA to produce acetoacetyl-CoA, a β -ketothiolase capable of condensing acetyl-CoA and propionyl-CoA to form β -ketovaleryl-CoA, e.g., BktB, a β -ketoacyl-CoA reductase capable of converting a β -ketoacyl-CoA to its corresponding β -hydroxyacyl-CoA, e.g., PhbB, and a PHA synthase capable
5 of producing P(3HB-co-3HV) copolymer from the β -hydroxyacyl-CoAs, e.g., PhbC, or combinations thereof, can be introduced into these organisms. Examples of such organisms include members of the genus *Clostridium*, such as *Clostridium arcticum*, *Clostridium novyi*, *Clostridium propionicum*, as well as *Megasphaera elsdenii*.

10

In addition, organisms that are capable of producing butyrate or butyryl-CoA from simple precursors are useful for producing C4/C6 copolymer, i.e., P(3HB-co-3HC) copolymer. For example, DNAs encoding a β -ketothiolase capable of condensing two molecules of acetyl-CoA to produce acetoacetyl-CoA,
15 a β -ketothiolase capable of condensing acetyl-CoA and butyryl-CoA to form β -ketocaproyl-CoA, e.g., BktB, a β -ketoacyl-CoA reductase capable of converting a β -ketoacyl-CoA to its corresponding β -hydroxyacyl-CoA, e.g., PhbB, and a PHA synthase capable of producing P(3HB-co-3HC) copolymer from the β -hydroxyacyl-CoA, e.g., *Nocardia corallina* PHA synthase, or
20 combinations thereof, can be introduced into these organisms. Examples of such organisms include species of the genera *Butyrivibrio*, *Clostridium* (e.g., *Clostridium populeti*), *Eubacterium*, and *Fusarium* (Patel and Agnew, 1988).

Plant Vectors

25

In plants, transformation vectors capable of introducing encoding DNAs involved in PHA biosynthesis are easily designed, and generally contain one or more DNA coding sequences of interest under the transcriptional control of 5'

and 3' regulatory sequences. Such vectors generally comprise, operatively linked in sequence in the 5' to 3' direction, a promoter sequence that directs the transcription of a downstream heterologous structural DNA in a plant; optionally, a 5' non-translated leader sequence; a nucleotide sequence that encodes a protein of interest; and a 3' non-translated region that encodes a polyadenylation signal which functions in plant cells to cause the termination of transcription and the addition of polyadenylate nucleotides to the 3' end of the mRNA encoding said protein. Plant transformation vectors also generally contain a selectable marker. Typical 5'-3' regulatory sequences include a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal. Vectors for plant transformation have been reviewed in Rodriguez et al. (1988), Glick et al. (1993), and Croy (1993).

Plant Promoters

Plant promoter sequences can be constitutive or inducible, environmentally- or developmentally-regulated, or cell- or tissue-specific. Often-used constitutive promoters include the CaMV 35S promoter (Odell et al., 1985), the enhanced CaMV 35S promoter, the Figwort Mosaic Virus (FMV) promoter (Richins et al., 1987), the mannopine synthase (*mas*) promoter, the nopaline synthase (*nos*) promoter, and the octopine synthase (*ocs*) promoter. Useful inducible promoters include heat-shock promoters (Ou-Lee et al., 1986; Ainley et al., 1990), a nitrate-inducible promoter derived from the spinach nitrite reductase gene (Back et al., 1991), hormone-inducible promoters (Yamaguchi-Shinozaki et al., 1990; Kares et al., 1990), and light-inducible promoters associated with the small subunit of RuBP carboxylase and LHCP gene families (Kuhlemeier et al., 1989; Feinbaum et al., 1991; Weisshaar et al.,

1991; Lam and Chua, 1990; Castresana et al., 1988; Schulze-Lefert et al., 1989). Examples of useful tissue-specific, developmentally-regulated promoters include the β -conglycinin 7S promoter (Doyle et al., 1986; Slighton and Beachy, 1987), and seed-specific promoters (Knutzon et al., 1992; Bustos et al., 1991; Lam and Chua, 1991; Stayton et al., 1991). Plant functional promoters useful for preferential expression in seed plastids include those from plant storage protein genes and from genes involved in fatty acid biosynthesis in oilseeds. Examples of such promoters include the 5' regulatory regions from such genes as napin (Kridl et al., 1991), phaseolin, zein, soybean trypsin inhibitor, ACP, stearyl-ACP desaturase, and oleosin. Seed-specific gene regulation is discussed in EP 0 255 378. Promoter hybrids can also be constructed to enhance transcriptional activity (Hoffman, U.S. Patent No. 5,106,739), or to combine desired transcriptional activity and tissue specificity.

15 A factor to be considered in the choice of promoters is the timing of availability of the necessary substrates during expression of the PHA biosynthetic enzymes. For example, if P(3HB-co-3HV) copolymer is produced in seeds from threonine, the timing of threonine biosynthesis and the amount of free threonine are important considerations. Karchi et al. (1994) have reported that threonine biosynthesis occurs rather late in seed development, similar to the timing of seed storage protein accumulation. For example, if enzymes involved in P(3HB-co-3HV) copolymer biosynthesis are expressed from the 7S seed-specific promoter, the timing of expression thereof will be concurrent with threonine accumulation.

25

Plant Transformation and Regeneration

A variety of different methods can be employed to introduce such vectors into plant protoplasts, cells, callus tissue, leaf discs, meristems, etc., to generate transgenic plants, including *Agrobacterium*-mediated transformation, particle gun delivery, microinjection, electroporation, polyethylene glycol-

mediated protoplast transformation, liposome-mediated transformation, etc. (reviewed in Potrykus, 1991). In general, transgenic plants comprising cells containing and expressing DNAs encoding enzymes facilitating PHA biosynthesis can be produced by transforming plant cells with a DNA

- 5 construct as described above via any of the foregoing methods; selecting plant cells that have been transformed on a selective medium; regenerating plant cells that have been transformed to produce differentiated plants; and selecting a transformed plant which expresses the enzyme-encoding nucleotide sequence.

- 10 Constitutive overexpression of, for example, a deregulated threonine deaminase employing the e35S or FMV promoter might potentially starve plants of certain amino acids, especially those of the aspartate family. If such starvation occurs, the negative effects may be avoided by supplementing the growth and cultivation media employed in the transformation and regeneration
- 15 procedures with appropriate amino acids. By supplementing the transformation/regeneration media with aspartate family amino acids (aspartate, threonine, lysine, and methionine), the uptake of these amino acids into the plant can reduce any potential starvation effect caused by an overexpressed threonine deaminase. Supplementation of the media with such
- 20 amino acids might thereby prevent any negative selection, and therefore any adverse effect on transformation frequency, due to the overexpression of a deregulated threonine deaminase in the transformed plant.

- The encoding DNAs can be introduced either in a single transformation
- 25 event (all necessary DNAs present on the same vector), a co-transformation event (all necessary DNAs present on separate vectors that are introduced into plants or plant cells simultaneously), or by independent transformation events (all necessary DNAs present on separate vectors that are introduced into plants or plant cells independently). Traditional breeding methods can
- 30 subsequently be used to incorporate the entire pathway into a single plant. Successful production of the PHA polyhydroxybutyrate in cells of *Arabidopsis*

has been demonstrated by Poirier et al. (1992), and in plastids thereof by Nawrath et al. (1994).

Specific methods for transforming a wide variety of dicots and obtaining
5 transgenic plants are well documented in the literature (Gasser and Fraley, 1989; Fisk and Dandekar, 1993; Christou, 1994; and the references cited therein).

Successful transformation and plant regeneration have been achieved in
10 the monocots as follows: asparagus (*Asparagus officinalis*; Bytebier et al. 1987); barley (*Hordeum vulgare*; Wan and Lemaux 1994); maize (*Zea mays*; Rhodes et al., 1988; Gordon-Kamm et al., 1990; Fromm et al., 1990; Koziel et al., 1993); oats (*Avena sativa*; Somers et al., 1992); orchardgrass (*Dactylis glomerata*; Horn et al., 1988); rice (*Oryza sativa*, including indica and japonica
15 varieties; Toriyama et al., 1988; Zhang et al., 1988; Luo and Wu 1988; Zhang and Wu 1988; Christou et al., 1991); rye (*Secale cereale*; De la Pena et al., 1987); sorghum (*Sorghum bicolor*; Cassas et al. 1993); sugar cane (*Saccharum* spp.; Bower and Birch 1992); tall fescue (*Festuca arundinacea*; Wang et al. 1992); turfgrass (*Agrostis palustris*; Zhong et al., 1993); and wheat (*Triticum aestivum*;
20 Vasil et al. 1992; Weeks et al. 1993; Becker et al. 1994).

Host Plants

Particularly useful plants for PHA copolymer production include those
25 that produce carbon substrates which can be employed for PHA biosynthesis, including tobacco, wheat, potato, *Arabidopsis*, and high oil seed plants such as corn, soybean, canola, oil seed rape, sunflower, flax, and peanut. Polymers that can be produced in this manner include copolymers incorporating both short chain length and medium chain length monomers, such as P(3HB-co-3HV)
30 copolymer.

If the host plant of choice does not produce the requisite fatty acid substrates in sufficient quantities, it can be modified, for example by mutagenesis or genetic transformation, to block or modulate the glycerol ester and fatty acid biosynthesis or degradation pathways so that it accumulates the appropriate substrates for PHA production.

P(3HB-co-3HV) Copolymer Composition

The P(3HB-co-3HV) copolymers of the present invention can comprise about 75-99% 3HB and about 1-25% 3HV based on the total weight of the polymer. More preferably, P(3HB-co-3HV) copolymers of the present invention comprise about 85-99% 3HB and about 1-15% 3HV. Even more preferably, such copolymers comprise about 90-99% 3HB and about 1-10% 3HV. P(3HB-co-3HV) copolymers comprising about 4%, about 8%, and about 12% 3HV on a weight basis possess properties that have made them commercially attractive for particular applications. One skilled in the art can modify P(3HB-co-3HV) copolymers of the present invention by physical or chemical means to produce copolymer derivatives having desirable properties different from those of the plant- or microbially-produced copolymer.

20

Optimization of P(3HB-co-3HV) copolymer production by the methods discussed herein is expected to result in yields of copolymer in the range of from at least about 1% to at least about 20% of the fresh weight of the plant tissue, organ, or structure in which it is produced. In bacteria, yields in the range of from at least about 1% to at least about 90% of cell dry weight is expected.

25

Plastid Targeting of Expressed Enzymes for PHA Biosynthesis

PHA polymer can be produced in plants either by expression of the appropriate enzymes in the cytoplasm (Poirier et al., 1992) by the methods described above, or more preferably, in plastids, where higher levels of PHA

30

production can be achieved (Nawrath et al., 1994). As demonstrated by the latter group, targeting of β -ketothiolase, acetoacetyl-CoA reductase, and PHB synthase to plastids of *Arabidopsis thaliana* results in the accumulation of high levels of PHB in the plastids without any readily apparent deleterious effects on plant growth and seed production. As branched-chain amino acid biosynthesis occurs in plant plastids (Bryan, 1980; Galili, 1995), overexpression therein of plastid-targeted enzymes, including a deregulated form of threonine deaminase, is expected to facilitate the production of elevated levels of α -ketobutyrate and propionyl-CoA. The latter can be condensed with acetyl-CoA by β -ketothiolase to form 3-ketovaleryl-CoA, which can then be further metabolized by a β -ketoacyl-CoA reductase to 3-hydroxyvaleryl-CoA, the precursor of the C5 subunit of P(3HB-co-3HV) copolymer. As there is a high carbon flux through acetyl-CoA in plastids, especially in seeds of oil-accumulating plants such as oilseed rape (*Brassica napus*), canola (*Brassica rapa*, *Brassica campestris*, *Brassica carinata*, and *Brassica juncea*), soybean (*Glycine max*), flax (*Linum usitatissimum*), and sunflower (*Helianthus annuus*) for example, targeting of the gene products of desired encoding DNAs to leucoplasts of seeds, or transformation of seed leucoplasts and expression therein of these encoding DNAs, are attractive strategies for achieving high levels of PHA biosynthesis in plants.

All of the enzymes discussed herein can be modified for plastid targeting by employing plant cell nuclear transformation constructs wherein DNA coding sequences of interest are fused to any of the available transit peptide sequences capable of facilitating transport of the encoded enzymes into plant plastids (partially summarized in von Heijne et al., 1991), and driving expression by employing an appropriate promoter. The sequences that encode a transit peptide region can be obtained, for example, from plant nuclear-encoded plastid proteins, such as the small subunit (SSU) of ribulose biphosphate carboxylase, plant fatty acid biosynthesis related genes including

acyl carrier protein (ACP), stearyl-ACP desaturase, β -ketoacyl-ACP synthase and acyl-ACP thioesterase, or LHCPII genes. The encoding sequence for a transit peptide effective in transport to plastids can include all or a portion of the encoding sequence for a particular transit peptide, and may also contain

5 portions of the mature protein encoding sequence associated with a particular transit peptide. Numerous examples of transit peptides that can be used to deliver target proteins into plastids exist, and the particular transit peptide encoding sequences useful in the present invention are not critical as long as delivery into a plastid is obtained. Proteolytic processing within the plastid then

10 produces the mature enzyme. This technique has proven successful not only with enzymes involved in PHA synthesis (Nawrath et al., 1994), but also with neomycin phosphotransferase II (NPT-II) and CP4 EPSPS (Padgett et al., 1995), for example.

15 Of particular interest are transit peptide sequences derived from enzymes known to be imported into the leucoplasts of seeds. Examples of enzymes containing useful transit peptides include those related to lipid biosynthesis (e.g., subunits of the plastid-targeted dicot acetyl-CoA

20 carboxylase, biotin carboxylase, biotin carboxyl carrier protein, α -carboxy-transferase, plastid-targeted monocot multifunctional acetyl-CoA carboxylase (Mr, 220,000); plastidic subunits of the fatty acid synthase complex (e.g., acyl carrier protein (ACP), malonyl-ACP synthase, KASI, KASII, KASIII, etc.); steroyl-ACP desaturase; thioesterases (specific for short, medium, and long chain acyl ACP); plastid-targeted acyl transferases (e.g., glycerol-3-phosphate:

25 acyl transferase); enzymes involved in the biosynthesis of aspartate family amino acids; phytoene synthase; gibberellic acid biosynthesis (e.g., *ent*-kaurene synthases 1 and 2); sterol biosynthesis (e.g., hydroxy methyl glutaryl-coA reductase); and carotenoid biosynthesis (e.g., lycopene synthase).

30 Exact translational fusions to the transit peptide of interest may not be optimal for protein import into the plastid. By creating translational fusions of

any of the enzymes discussed herein to the precursor form of a naturally imported protein or C-terminal deletions thereof, one would expect that such translational fusions would aid in the uptake of the engineered precursor protein into the plastid. For example, Nawrath et al., (1994) used a similar
5 approach to create the vectors employed to introduce the PHB biosynthesis genes of *A. eutrophus* into *Arabidopsis*.

It is therefore fully expected that targeting of the enzymes discussed in Examples 1-8 to leaf chloroplasts or seed plastids such as leucoplasts by
10 fusing transit peptide gene sequences thereto will further enhance *in vivo* conditions for the biosynthesis of PHAs in plants.

Plastid Transformation for Expression of Enzymes Involved in PHA Biosynthesis

15 Alternatively, enzymes facilitating the biosynthesis of metabolites such as threonine, α -ketobutyrate, propionyl-CoA, 3-ketovaleryl-CoA, 3-hydroxyvaleryl-CoA, and PHAs discussed herein can be expressed *in situ* in plastids by direct transformation of these organelles with appropriate recombinant
20 expression constructs. Constructs and methods for stably transforming plastids of higher plants are well known in the art (Svab et al., 1990; Svab et al., 1993; Staub et al., 1993; Maliga et al., U.S. Patent No. 5,451,513; PCT International Publications WO 95/16783, WO 95/24492, and WO 95/24493). These methods generally rely on particle gun delivery of DNA containing a
25 selectable marker in addition to introduced DNA sequences for expression, and targeting of the DNA to the plastid genome through homologous recombination. Transformation of a wide variety of different monocots and dicots by particle gun bombardment is routine in the art (Hinchee et al., 1994; Walden and Wingender, 1995).

DNA constructs for plastid transformation generally comprise a targeting segment comprising flanking DNA sequences substantially homologous to a predetermined sequence of a plastid genome, which targeting segment enables insertion of DNA coding sequences of interest into the plastid genome by homologous recombination with said predetermined sequence; a selectable marker sequence, such as a sequence encoding a form of plastid 16S ribosomal RNA that is resistant to spectinomycin or streptomycin, or that encodes a protein which inactivates spectinomycin or streptomycin (such as the *aadA* gene), disposed within said targeting segment, wherein said selectable marker sequence confers a selectable phenotype upon plant cells, substantially all the plastids of which have been transformed with said DNA construct; and one or more DNA coding sequences of interest disposed within said targeting segment relative to said selectable marker sequence so as not to interfere with conferring of said selectable phenotype. In addition, plastid expression constructs also generally include a plastid promoter region and a transcription termination region capable of terminating transcription in a plant plastid, wherein said regions are operatively linked to the DNA coding sequences of interest.

A further refinement in chloroplast transformation/expression technology that facilitates control over the timing and tissue pattern of expression of introduced DNA coding sequences in plant plastid genomes has been described in PCT International Publication WO 95/16783. This method involves the introduction into plant cells of constructs for nuclear transformation that provide for the expression of a viral single subunit RNA polymerase and targeting of this polymerase into the plastids via fusion to a plastid transit peptide. Transformation of plastids with DNA constructs comprising a viral single subunit RNA polymerase-specific promoter specific to the RNA polymerase expressed from the nuclear expression constructs operably linked to DNA coding sequences of interest permits control of the plastid expression constructs in a tissue and/or developmental specific manner in plants comprising both the nuclear polymerase construct and the plastid

expression constructs. Expression of the nuclear RNA polymerase coding sequence can be placed under the control of either a constitutive promoter, or a tissue- or developmental stage-specific promoter, thereby extending this control to the plastid expression construct responsive to the plastid-targeted, nuclear-
5 encoded viral RNA polymerase. The introduced DNA coding sequence can be a single encoding region, or may contain a number of consecutive encoding sequences to be expressed as an engineered or synthetic operon. The latter is especially attractive where, as in the present invention, it is desired to introduce multigene biochemical pathways into plastids. This approach is not
10 practical using standard nuclear transformation techniques since each gene introduced therein must be engineered as a monocistron, including an encoded transit peptide and appropriate promoter and terminator signals. Individual gene expression levels may vary widely among different cistrons, thereby possibly adversely affecting the overall biosynthetic process. This can be
15 avoided by the chloroplast transformation approach.

Production of Transgenic Plants Comprising Genes for PHA Biosynthesis

20

Plant transformation vectors capable of delivering DNAs (genomic DNAs, plasmid DNAs, cDNAs, or synthetic DNAs) encoding PHA biosynthetic enzymes and other enzymes for optimizing substrate pools for PHA biosynthesis as discussed above in Examples 1-8 can be easily designed.
25 Various strategies can be employed to introduce these encoding DNAs to produce transgenic plants capable of biosynthesizing high levels of PHAs, including:

1. Transforming individual plants with an encoding DNA of interest.
- 30 Two or more transgenic plants, each containing one of these DNAs, can then be grown and cross-pollinated so as to produce hybrid plants containing the two

DNAs. The hybrid can then be crossed with the remaining transgenic plants in order to obtain a hybrid plant containing all DNAs of interest within its genome.

2. Sequentially transforming plants with plasmids containing each of the
5 encoding DNAs of interest, respectively.

3. Simultaneously cotransforming plants with plasmids containing each of the encoding DNAs, respectively.

10 4. Transforming plants with a single plasmid containing two or more encoding DNAs of interest.

5. Transforming plants by a combination of any of the foregoing techniques in order to obtain a plant that expresses a desired combination of
15 encoding DNAs of interest.

Traditional breeding of transformed plants produced according to any one of the foregoing methods by successive rounds of crossing can then be carried out to incorporate all the desired encoding DNAs in a single homozygous plant
20 line (Nawrath et al., 1994; PCT International Publication WO 93/02187). Similar strategies can be employed to produce bacterial host cells engineered for optimal PHA production.

In methods 2 and 3, the use of vectors containing different selectable
25 marker genes to facilitate selection of plants containing two or more different encoding DNAs is advantageous. Examples of useful selectable marker genes include those conferring resistance to kanamycin, hygromycin, sulphonamides, glyphosate, bialaphos, and phosphinothricin.

30 **Stability of Transgene Expression**

As several overexpressed enzymes may be required to produce optimal levels of substrates for copolymer formation, the phenomenon of co-suppression may influence transgene expression in transformed plants. Several strategies can be employed to avoid this potential problem (Finnegan
5 and McElroy, 1994).

One commonly employed approach is to select and/or screen for transgenic plants that contain a single intact copy of the transgene or other encoding DNA (Assaad et al., 1993; Vaucheret, 1993; McElroy and Brettell,
10 1994). *Agrobacterium*-mediated transformation technologies are preferred in this regard.

Inclusion of nuclear scaffold or matrix attachment regions (MAR) flanking a transgene has been shown to increase the level and reduce the
15 variability associated with transgene expression in plants (Stief et al., 1989; Breyne et al., 1992; Allen et al., 1993; Mlynarova et al., 1994; Spiker and Thompson, 1996). Flanking a transgene or other encoding DNA with MAR elements may overcome problems associated with differential base
composition between such transgenes or encoding DNAs and integrations sites,
20 and/or the detrimental effects of sequences adjacent to transgene integration sites.

The use of enhancers from tissue-specific or developmentally-regulated genes may ensure that expression of a linked transgene or other encoding DNA
25 occurs in the appropriately regulated manner.

The use of different combinations of promoters, plastid targeting sequences, and selectable markers for introduced transgenes or other encoding DNAs can avoid potential problems due to *trans*-inactivation in cases where
30 pyramiding of different transgenes within a single plant is desired.

5 the same gene, but with altered expression characteristics, should obviate this problem (Yoder and Goldsborough, 1994).

10 the present invention.

15 would be obvious to one skilled in the art are intended to be included within the
scope of the following claims.

References Cited

Ainley et al., (1990) *Plant Mol. Biol.* 14: 949.

5 Allen et al., (1993) *Plant Cell* 5: 603.

Altschul et al., (1990) *J. Mol. Biol.* 215: 403.

Assaad et al., (1993) *Plant Mol. Biol.* 22: 1067.

10

Ausubel et al., (1989) *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc.

Back et al., (1991) *Plant Mol. Biol.* 17: 9.

15

Becker et al., (1994) *Plant J.* 5: 299.

Bisswanger, (1981) *J. Biol. Chem.* 256: 815.

20 Bower and Birch, (1992) *Plant J.* 2: 409.

Boy et al., (1979) *Biochimie* 61: 1151.

Boylan and Dekker, (1981) *J. Biol. Chem.* 256: 1809.

25

Bradford, (1976) *Anal. Biochem.* 72: 248.

Brandl et al., (1990) *Adv. Biochem. Eng. Biotech.* 41: 77.

30 Brandl et al., (1989) *Int. J. Biol. Macromol.* 11: 49.

"FBI" 03-16-00

-

- 30

Feldmann et al., (1994) In: *Arabidopsis*, Meyerowitz and Somerville (Eds.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., p. 137.

- 5 Finnegan and McElroy, (1994) *Bio/Technology* 12: 883.

Fisher and Eisenstein, (1993) *J. Bacteriol.* 175: 6605.

Fisk and Dandekar, (1993) *Scientia Horticulturae* 55: 5.

10

Fosket, (1994) *Plant Growth and Development*, Academic Press, Inc., San Diego, p. 132.

Fraley et al., (1983) *Proc. Natl. Acad. Sci USA.* 80: 4803.

15

Frankard et al., (1991) *Theor. Appl. Genet.* 82: 273.

Fries, (1962) *Physiol. Plant.* 15: 566.

- 20 Fromm et al., (1990) *Bio/Technology* 8: 833.

Fry et al., (1987) *Plant Cell Reports* 6: 321.

Galili, (1995) *Plant Cell* 7: 899.

25

Gasser and Fraley, (1989) *Science* 244: 1293.

Glick et al., (1993) *Methods in Plant Molecular Biology and Biotechnology*, CRC Press, Boca Raton, Fla.

30

Gordon-Kamm et al., (1990) *Plant Cell* 2: 603.

Goss and Datta, (1985) *Mol. Gen. Genet.* 201: 308.

Gruys et al., (1989) *Biochemistry* 28: 9071-9080.

5

Halarnkar et al., (1988) *Phytochemistry* 27: 997.

Halarnkar and Blomquist, (1989) *Comp. Biochem. Physiol.* 92: 227.

10 Haywood et al., (1988a) *FEMS Microbiol. Lett.* 52: 259.

Haywood et al., (1988b) *FEMS Microbiol. Lett.* 52: 91.

Haywood, (1989) *FEMS Microbiol Lett.* 57: 1.

15

Hibberd et al., (1980) *Planta* 148: 183.

Hinchee et al., (1988) *Bio/Technology* 6: 915.

20 Hinchee et al., (1994) In: *Plant Cell and Tissue Culture*, I. Vasil and T. Thorpe (Eds.), Kluwer Academic Publishers, Netherlands, p. 231.

Hoffman, U.S. Patent No. 5,106,739

25 Horn et al., (1988) *Plant Cell Rep.* 7: 469.

Horsch et al., (1985) *Science* 227: 1229.

Hunkapiller et al., (1983) *Methods Enzymol.* 91: 399.

30

Jan et al., (1995) *Analytical Biochemistry* 225: 258.

- Jetten et al., (1995) *Appl. Microbiol. Biotechnol.* 43: 76.
- Jetten and Sinskey, (1995) *Critical Rev. Biotechnol.* 15: 73.
- 5 Karchi et al., (1993) *Plant J.* 3: 721.
- Kares et al., (1990) *Plant Mol. Biol.* 15: 905.
- 10 Katake and Ono, (1964) *Shokukin Eiseigaku Zasshi* 5: 39.
- Kidwell et al., (1995) *Appl. Environ. Microbiol.* 61: 1391.
- Kim and Patel, (1992) *J. Biol. Chem.* 267: 5128.
- 15 Knutzon et al., (1992) *Proc. Natl. Acad. Sci USA* 89: 2624.
- Koncz and Shell, (1986) *Mol. Gen. Genet.* 204: 283.
- 20 Kovach, M. et al. (1994) *Biotechniques.* 16:800.
- Kozziel et al., (1993) *Bio/Technology* 11: 194.
- Kridl et al., (1991) *Seed Sci. Res.* 1: 209.
- 25 Kuhlemeier et al., (1989) *Plant Cell* 1: 471.
- Lam and Chua, (1990) *Science* 248: 471.
- 30 Lam and Chua, (1991) *J. Biol. Chem.* 266: 17131

LaRossa, (1987) *J. Bacteriol.* 169: 1372.

Laties and Hoell, (1967) *Phytochemistry* 6: 49.

Lawther et al., (1987) *Nucl. Acids Res.* 15: 2137.

5

Layman, (Oct. 31, 1994) *Chem. & Eng. News*, p. 10.

Lee et al., (1995) *J. Fermentation and Bioengineering* 79: 177.

10

Lemoigne, (1926) *Bull. Soc. Chim. Biol.* (Paris) 8: 770.

Lindsay, (1992) *Modern Plastics* 2: 62.

Lowe et al., (1983) *Biochem. J.* 215: 133.

15

Luo and Wu, (1988) *Plant Mol. Biol. Rep.* 6: 165.

Maliga et al., U.S. Patent No. 5,451,513.

20

Maliga et al., PCT International Publication WO 95/16783.

Maliga et al., PCT International Publication WO 95/24492.

Maliga et al., PCT International Publication WO 95/24493.

25

Manchak and Page, (1994) *Microbiology* 140: 953.

Marcus and Dekker, (1993) *Biochimica et Biophysica Acta* 1164: 299.

30

Masamune et al., (1989) *Pure & Appl. Chem.* 6: 303.

McElroy and Brettell, (1994) *TIBTECH* 12: 62.

Miller and Schulz, (1993) *Science* 259: 965.

Möckel et al., (1994) *Molec. Microbiol.* 13: 833.

5

Monod et al., (1963) *J. Mol. Biol.* 6: 306.

Morelli et al., (1985) *Nature* 315: 200.

10

Motoyama et al., (1994) *Appl. Microbiol. Biotech.* 42: 67.

Mourad and King, (1995) *Plant Physiol.* 107: 43

Mlynarova et al., (1994) *Plant Cell* 6: 417.

15

Muhitch, (1995) *Plant Physiol.* 108 (2 Suppl.): 71.

Murashige and Skoog, (1962) *Physiol. Plant.* 15: 473.

20

Nakamura et al., (1992) *Int. J. Biol. Macromol.* 14: 321.

Nakamura et al., (1991) *Macromol. Rep.* A28 (Suppl. 1): 15.

Nawrath et al., (1994) *Proc. Natl. Acad. Sci USA* 91: 12760.

25

Odell et al., (1985) *Nature* 313: 810.

Ou-Lee et al., (1986) *Proc. Natl. Acad. Sci USA* 83: 6815.

- Padgette et al., (1996) In: *Herbicide Resistant Crops: Agricultural, Environmental, Economic, Regulatory, and Technical Aspects*. S. O. Duke (Ed.), CRC Press, N.Y., p. 53.
- 5 Padgette et al., (1995) *Crop Sci.* 35: 1451.
Patel and Walt, (1987) *J. Biol. Chem.* 262: 7132.
- Patel and Agnew, (1988) *Arch. Microbiol.* 150:267-271.
- 10 Peoples and Sinskey, (1989a) *J. Biol. Chem.* 264: 15293.
Peoples and Sinskey, (1989b) *J. Biol. Chem.* 264: 15298.
- Petrosyan et al., (1969) *Vopr. Mikrobiol.* 181.
- 15 Pettit et al., (1978) *Proc. Nat. Acad. Sci. USA* 75: 4881.
- Poirier et al., (1992) *Science* 256: 520.
- 20 Poirier et al., (1995) *Bio/Technology* 13: 142.
- Poston, (1977) *Science* 195: 301.
- Poston., (1978) *Phytochemistry* 17: 401.
- 25 Potrykus, (1991) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 42: 205.
- Priefert and Steinbüchel (1992) *J. Bact.* 174: 6590
- 30 Qureshi et al., (1982) *J. Chrom.* 249: 333.

Radke et al., (1988) *Theor. Appl. Genet.* 75: 685.

Register et al., (1994) *Plant Mol. Biol.* 25: 951.

5 Reed and Willms (1966) *Methods Enzymol.* 9: 247-265.

Rhodes et al., (1988) *Science* 240: 204.

Richins et al., (1987) *NAR* 20: 8451.

10 Rodriguez et al., (1988) *Vectors: A Survey of Molecular Cloning Vectors and Their Uses*, Butterworths, Boston.

Sambrook et al., (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

15

Schubert et al., (1988) *J. Bacteriol.* 170: 5837.

Schulz (1974) *J. Biol. Chem.* 249: 2704.

20 Schulze et al., (1992) *Eur. J. Biochem.* 206: 427.

Schulze et al., (1993) *Eur. J. Biochem.* 211: 591.

Schulze-Lefert et al., (1989) *EMBO J.* 8: 651.

25

Shah et al., (1986) *Science* 233: 478.

Shaner and Singh, (1993) *Plant Physiol* 103: 1221.

30 Shaul and Galili, (1992a) *Plant Physiol.* 100: 1157.

13618-21-38

Shaul and Galili, (1992b) *Plant J.* 2: 203.

Shaul and Galili, (1993) *Plant Mol. Biol.* 23: 759.

5 Shizuta and Tokushige, (1971) *Methods Enzymol.* 17: 575.

Shizuta and Hayashi, (1975) *Curr. Topics Cell. Reg.* 11: 99.

Slater et al., (1988). *J. Bacteriol.* 170: 4431.

10

Slater et al., (1992) *Appl. Environ. Microbiol.* 58: 1089.

Slighton and Beachy, (1987) *Planta* 172: 356.

15 Somers et al., (1992) *Bio/Technology* 10: 1589.

Speckhard and Frey, (1975) *Biochem. Biophys. Res. Commun.* 62: 614-620.

Spiker and Thompson, (1996) *Plant Physiol.* 110: 15.

20

Staba, E. J., (1969) *Recent Advances in Phytochemistry*, Vol. 2, Seikel et al. (Eds.), Appleton Century Crofts, N.Y.

Stark et al., (1992) *Science* 258: 287.

25

Staub and Maliga, (1993) *EMBO J.* 12: 601.

Stayton et al., (1991) *Aust. J. Plant. Physiol.* 18: 507.

30 Stief et al., (1989) *Nature* 341: 343.

Steinbüchel, (1991) In: *Biomaterials*, D. Byrom (Ed.), MacMillan, Basingstoke, pp. 123-213.

Steinbüchel and Pieper, (1992) *Appl. Microbiol. Biotechnol.* 37: 1.

5

Steginsky et al., (1985) *J. Biol. Chem.* 260: 13690.

Strauss et al., (1985) *Planta* 163: 554.

Svab and Maliga, (1993) *Proc. Natl. Acad. Sci. USA* 90: 913.

10

Svab et al., (1990) *Proc. Natl. Acad. Sci. USA* 87: 8526.

Taillon et al., (1988) *Gene* 63: 245.

15 Tholozan et al., (1992) *Arch. Microbiol.* 157: 249.

Tholozan et al., (1994) *Arch. Microbiol.* 162: 401.

Tombolini et al., (1995) *Microbiology* 141: 2553.

20

Toriyama et al., (1988) *Bio/Technology* 6: 10.

Van den Broeck et al., (1985) *Nature* 313: 358.

25 Van Dyk and LaRossa (1987) *Mol. Gen. Genet.* 207: 435.

Vasil et al., (1992) *Bio/Technology* 10:667

Vaucheret, (1993) *C.R. Acad. Sci. Paris, Science de la vie / Life Sciences* 316:

30 1471.

Von Heijne et al., (1991) *Plant Mol. Biol. Rep.* 9: 104.

Walker (1987) In: *Eschericia coli and Salmonella typhimurium: Cellular and Molecular Biology*, F.C. Neidhart et al. (Eds.), American Society for

5 Microbiology, Washington, D.C., pp. 13346-1357.

Weisshaar et al., (1991) *EMBO J.* 10: 1777.

Williams et al., (1994) *Appl. Microbiol. Biotechnol.* 40: 717.

10

Walden and Wingender, (1995) *TIBS* 13: 324.

Wallis, (1994) *J. Mol. Biol.* 236: 209.

15 Wan and Lemaux, (1994) *Plant Physiol.* 104: 37

Wang et al., (1992) *Bio/Technology* 10: 691.

Weeks et al., (1993) *Plant Physiol.* 102: 1077.

20

Yamaguchi-Shinozaki et al., (1990) *Plant Mol. Biol.* 15: 905.

Yoder and Goldsbrough, (1994) *Bio/Technol.* 12: 263-267.

25 Yoon et al., (1995) *J. Ferment. Bioeng.* 80: 350

Zhang et al., (1994) *Appl. Environ. Microbiol.* 60: 1198-1205.

Zhang and Wu, (1988) *Theor. Appl. Genet.* 76: 835.

30

Zhang et al., (1988) *Plant Cell Rep.* 7: 379.

Zhong et al., (1993) *Plant Cell Rep.* 13: 1.

Zinszner et al., (1994) *Genes Dev.* 8: 2513.

1-03693-1-03693-1